

Combinatorial Stability of Nondeterministic Systems

P. Duarte¹, M. J. Torres²

¹ *CMAF, Faculdade de Ciências da Universidade de Lisboa, pduarte@ptmat.fc.ul.pt, with [partial] support from the FCT, program POCTI (Portugal/FEDER-EU)*

² *Departamento de Matemática da Universidade do Minho, jtorres@math.uminho.pt, with [partial] support from CMAT (Centro de Matemática da Universidade do Minho) and FCT, program POCTI (Portugal/FEDER-EU)*

Abstract: *We introduce and study, from a combinatorial-topological viewpoint, some semigroups of continuous non-deterministic dynamical systems. Combinatorial stability, i.e. the attractors' combinatorics persistence, is characterized and its genericity established. Some implications on topological (deterministic) dynamics are drawn.*

1 Introduction

In the framework of smooth deterministic dynamical systems, i.e. smooth maps, diffeomorphisms, and flows on finite dimensional compact manifolds, it has been long believed that *typical dynamical systems should exhibit a finite number of attractors, their basins of attraction covering almost everything.*

S. Smale program in the early sixties aimed proving the (topological) genericity of structurally stable systems. The concept of structural stability was introduced in the thirties by Andronov-Pontryagin. It means that under small perturbations the dynamics is topologically equivalent to the original system. Although Smale's program was proved to be wrong one decade later, it played a fundamental role in the development of dynamical systems' theory. It led to the construction of Hyperbolic Theory, studying uniform hyperbolicity and characterizing structural stability as being essentially equivalent to uniform hyperbolicity. Of course the scenario above is true for hyperbolic (Axiom A) systems.

More recently J. Palis conjectured, see [P], that every dynamical system can be approximated by one with finitely many attractors, each having a stochastically stable physical measure, whose basins of attraction cover almost every point in state space. A *physical measure* of an attractor is one that describes the system time average for a positive volume set of initial states. The set of all such initial states is called the physical measure basin of attraction. A measure on an attractor is said to be *stochastically stable* if it is stable under small stochastic perturbations of the deterministic system. More precisely, introducing a random noise, the limit measures of the random perturbations approach the attractor physical measure as the noise level tends to zero. The key idea of introducing a random noise in a deterministic system, and then looking at the limit measures as the noise level tends to zero, is usually attributed to Kolmogorov. Under very mild conditions, a random noise can have a powerful simplifying effect on the dynamics complexity of a deterministic system. Namely, see for instance [A], under arbitrary small random perturbations any deterministic system has finitely many attractors. Another simplifying effect may be seen on the Perron-Frobenius operator, whose fixed points are precisely the system invariant measures. In general the spectrum of this linear operator, reflecting the action of dynamics upon measures supported in state space, can be complex, but when a random noise is turned-on this usually makes the operator compact or quasi-compact with pure point spectrum. A compact operator can be, spectrally speaking, well approximated by finite dimension operators. Thus random perturbations of a deterministic system may, just as well, be considered on finite (discrete) approximations of state space. Finite state Markov chains are the stochastic, or random dynamical systems on a finite state space. One may think that these dynamical systems are what we actually see when running computer simulations of deterministic dynamical systems. Each such dynamical system is specified by a *stochastic matrix* with the state probability transitions. The stochastic matrix is the Perron operator of this finite state system. The Markov chain also determines an *oriented graph*, encapsulating some qualitative aspects of the system behaviour. The Theory of finite state Markov Chains, establishes a correspondence between spectral properties of the stochastic matrix on one side, and combinatorial properties of the corresponding graph on the other hand.

This theory can be nicely extended into a theory for continuous, or lower semi-continuous, random dynamical systems of Markov type on a compact manifold X . In the present article we develop such a theory in its topological and combinatorial aspects. Then in a complementary paper we shall address the correspondent spectral

theory and its relation with the results here obtained. The main novelty with respect to finite state Markov chain' s theory is that in this context, because we are dealing with continuous systems, it makes sense defining stability: combinatorial stability or spectral stability. The core of the present article will be to establish and characterize combinatorial stability.

We shall call *nondeterministic dynamical system* on a state space X to any multi-valued mapping $\varphi : X \rightarrow \mathcal{P}(X)$, where $\mathcal{P}(X)$ denotes the set of all X subsets. Here the state space X will be a compact manifold and we make a fundamental assumption on the state transition sets $\varphi(x)$ ($x \in X$). They are always non-empty connected open sets. The lower semi-continuity of a nondeterministic dynamical system φ w.r.t. the Hausdorff distance is then equivalent to the openness of $graph(\varphi)$. Such systems will be referred throughout this work as *open maps*. In section 3 we define several topological spaces of open sets. Then in section 4 we introduce and topologize some semigroups of open maps. Any open map φ can be identified with its graph $graph(\varphi)$ and, therefore, seen as an open set in $X \times X$. Thus, semigroups of open maps can be given topologies as subspaces of topological spaces of open sets. The key concept of *topological semigroup of open maps*, to which the main theorems 5.3 and 5.4 apply, is defined here. Examples of topological semigroups of open maps are given. In section 5 we define *recurrency* and *chain-recurrency*. Briefly, given any open map φ , a sequence x_0, x_1, \dots, x_n such that $x_i \in \varphi(x_{i-1})$ for all $i = 1, \dots, n$ will be called a φ -*orbit*, and we shall say that x_n is a φ -*iterate* of x_0 . If for every $\epsilon > 0$, y is an φ_ϵ^* -iterate of x , where φ_ϵ^* is the open map whose graph is ϵ -radius ball of $graph(\varphi)$, we say that y is a φ -*pseudo iterate* of x . The *recurrent* and *chain-recurrent sets* of φ are defined respectively by $\Omega(\varphi) = \{x \in X : x \text{ is a } \varphi\text{-iterate of } x\}$ and $R(\varphi) = \{x \in X : x \text{ is a } \varphi\text{-pseudo iterate of } x\}$. Both these sets split into equivalence classes, each class being formed by states which are accessible from each other. The set of all these classes is then partially ordered by the dynamics of φ . At the bottom of this hierarchy are two special limit sets: the *final recurrent* and the *final chain-recurrent sets*, denoted respectively by $\Omega_{\text{final}}(\varphi)$ and $R_{\text{final}}(\varphi)$, of all states $x \in \Omega(\varphi)$ ($x \in R_{\text{final}}(\varphi)$) such that every iterate (pseudo-iterate) of x still has some iterate (pseudo-iterate) which comes back to x . These limit sets contain all the asymptotic dynamical behaviour of φ . They both decompose into a finite number of connected pieces which are permuted by φ . The restriction of φ to each of these pieces being in some sense irreducible. See theorems 5.1 and 5.2. We say that two open maps φ and ψ are *combinatorially equivalent* iff there is a bijective conjugation between both maps' actions on the final recurrent sets' connected components. Given

a topological semigroup of open maps \mathcal{O}_1 , an open map $\varphi \in \mathcal{O}_1$ will be called \mathcal{O}_1 -combinatorially stable when there is a neighbourhood of φ in \mathcal{O}_1 where all open maps are combinatorially equivalent to φ . Since any iterate is also a pseudo iterate, there is a natural relation between the connected pieces of $\Omega_{\text{final}}(\varphi)$ and those of $R_{\text{final}}(\varphi)$. In theorem 5.3 we characterize combinatorial stability: for any topological semigroup of open maps \mathcal{O}_1 this relation is bijective *if and only if* φ is \mathcal{O}_1 -combinatorially stable. We also prove the genericity of combinatorial stability for topological semigroups of open maps, see theorem 5.4. In the end of section 5 we characterize the continuity of the attractors, i.e. the connected components of $\Omega_{\text{final}}(\varphi)$ or $R_{\text{final}}(\varphi)$, see theorem 5.5. When a combinatorially stable map is a continuity point of the attractors we say that it is *topologically stable*. Finally we establish, see theorem 5.6, that (in a topological sense) any generic open map φ , in some topological semigroup of open maps \mathcal{O}_1 , is topologically stable. In the last section, section 6, we derive some applications to the deterministic dynamics of continuous mappings. We shall denote by $\mathcal{C}^0(X)$ the space of all continuous mappings $f : X \rightarrow X$ with the usual C^0 topology of uniform convergence. Given any map $f \in \mathcal{C}^0(X)$ we define chain-recurrency and the final chain-recurrent set, $R_{\text{final}}(f)$, in a similar way as for open maps. Let $\mathcal{S} \subseteq \mathcal{C}^0(X)$ be a topological space or semigroup such that the inclusion $\mathcal{S} \hookrightarrow \mathcal{C}^0(X)$ is continuous. Let us say that $f \in \mathcal{S}$ is *strongly combinatorially stable* in \mathcal{S} iff there is a neighbourhood of f in \mathcal{S} where all maps are combinatorially equivalent to f , in the sense that there is a bijective conjugation between both maps' actions on the final chain-recurrent sets' connected components. If $\mathcal{S} = \mathcal{C}^0(X)$, this concept becomes too strong: there is no strongly combinatorially stable map in $\mathcal{C}^0(X)$, but in general with some smoothness assumption on \mathcal{S} , the set of all strongly combinatorially stable maps in \mathcal{S} is a non trivial open subset of \mathcal{S} . In general this open set is not dense in \mathcal{S} , as Newhouse examples with infinitely many sinks show. For instance, if $\mathcal{S} = \mathcal{C}^1(X)$ is the space of class C^1 mappings $f : X \rightarrow X$, systems with finitely many robustly transitive attractors (Axiom A in particular) are strongly combinatorially stable. Strong stability will not be addressed in the present article. Instead we shall work with a weaker concept. Given an open map φ , if $\text{graph}(f) \subseteq \text{graph}(\varphi)$, we write $f \prec \varphi$ and say that f is subordinated to φ . We say that f is *weakly combinatorially stable* if for every sufficiently small perturbation φ of f , with $f \prec \varphi$, the combinatorics of their final chain-recurrent sets are the same. We shall call *attractor*, in this paper, to any connected invariant chain-transitive set $A \subseteq X$ which attracts a whole neighbourhood of A . In theorem 6.1 we show that a map $f \in \mathcal{C}^0(X)$ is weakly combinatorially stable iff f has finitely many attractors iff

$R_{\text{final}}(f)$ has finitely many connected components. We should emphasize that 'weak combinatorial stability' is a really feeble concept. For instance, all volume preserving maps are weakly combinatorially stable. By the Poincaré recurrence theorem, every volume preserving map is chain transitive and, therefore, has the whole manifold as its unique attractor. In theorem 6.2 we show that the set of maps $f \in \mathcal{C}^0(X)$ with finitely many attractors is dense in $\mathcal{C}^0(X)$. In view of Palis' conjecture, it would be much more interesting proving this theorem for spaces of smoother functions $\mathcal{S} \subseteq \mathcal{C}^r(X)$, with $r \geq 1$. Thus, in order to allow for possible generalizations we have proved an abstract result, see theorem 6.3, saying that the same conclusion (density of maps with finitely many attractors) holds in every space $\mathcal{S} \subseteq \mathcal{C}^0(X)$ having the *skeleton-chain perturbation property*. Of course we only prove that $\mathcal{S} = \mathcal{C}^0(X)$ satisfies this condition, but because this key property has a rather geometric nature, we hope in the future to be able to develop new perturbation techniques for proving that other spaces of smoother functions also share this property.

2 Notation

Throughout this work X will denote a compact Riemannian manifold of dimension n , d will be the geodesic distance on X and m will be the corresponding normalized ($m(X) = 1$) Riemannian volume. X will be the state space of all systems that we consider.

Given sets $U, V \subseteq X$ we shall use the following notation.

1. $B_r(U)$, respectively $\overline{B}_r(U)$, denotes the set of all points whose distance d to U is less, respectively less or equal, than r .
2. $\rho_H(U, V) = \inf \{ r > 0 : U \subseteq B_r(V) \text{ and } V \subseteq B_r(U) \}$, denotes the Hausdorff distance between U and V .
3. $\rho_m(U, V) = m(U \setminus V) + m(V \setminus U)$, stands for the L^1 distance between the functions I_U and I_V , where $I_U(x) = \begin{cases} 1 & \text{if } x \in U \\ 0 & \text{if } x \notin U \end{cases}$.
4. $H^{n-1}(U)$ is the $(n - 1)$ -dimensional Hausdorff measure of U .

5. \bar{U} is the closure of U in X .
6. U° is the interior of U in X .
7. ∂U is the topological boundary of U in X .
8. U^e is the exterior of U in X .
9. $\widehat{U} = (\bar{U})^\circ$ is the interior of U 's closure in X .
10. U^c is the complement of U in X .
11. $|\Xi|$ is the number of elements in a set Ξ .

Similar notations will be used on $X \times X$, where d will stand for the metric,

$$d((x_1, y_1), (x_2, y_2)) = \max\{d(x_1, x_2), d(y_1, y_2)\}.$$

Since we will work in a semigroup of maps without the identity element, we shall consider the set \mathbb{N} as the set of all positive integers.

3 Topological Spaces of Open Sets

Let $\mathcal{U}(X)$ denote the space of all non-empty connected open subsets of X , and $\mathbb{U}(X) = \mathcal{U}(X)/\sim$ be the quotient space by the equivalence relation $U \sim V \Leftrightarrow \bar{U} = \bar{V} \Leftrightarrow \widehat{U} = \widehat{V}$, where \widehat{U} denotes the interior of \bar{U} .

An open set $U \subseteq X$ is said to be *geodesically r -regular*, or simply *r -regular*, if it is a union of open geodesic balls of radius $\geq r$. Note that, since any ball of radius $> r$ is itself a union of balls of radius r , any r -regular open set U is a union of balls of radius r . We shall denote by $\mathcal{U}^r(X)$ the set of all non-empty connected r -regular open sets. For $r = 0$ we set $\mathcal{U}^0(X) = \bigcup_{r>0} \mathcal{U}^r(X)$. Then, accordingly, the quotients of these spaces will be denoted by $\mathbb{U}^r(X) = \mathcal{U}^r(X)/\sim$, for every $r \geq 0$.

We are going to consider several topologies on the spaces $\mathcal{U}(X)$, $\mathcal{U}^r(X)$, $\mathbb{U}(X)$ and $\mathbb{U}^r(X)$. In the next section we shall deal with topological semigroups of *open maps*, which will be realized as subspaces of $\mathcal{U}(X \times X)$ and $\mathbb{U}(X \times X)$. Let us now give some abstract definitions:

Definition 3.1. We say that a family $\{U_\epsilon^*\}_{\epsilon>0}$, of open connected sets, is an outer approximation of U , if and only if

1. $\bar{U} = \bigcap_{\epsilon>0} U_\epsilon^*$, and
2. for all ϵ_1, ϵ_2 , $\epsilon_1 > \epsilon_2 > 0 \Rightarrow \bar{U}_{\epsilon_2}^* \subseteq U_{\epsilon_1}^*$.

Analogously, we say that a family $\{U_\epsilon^\circ\}_{\epsilon>0}$, of open connected sets, is an inner approximation of U , if and only if

1. $U = \bigcup_{\epsilon>0} U_\epsilon^\circ$, and
2. for all ϵ_1, ϵ_2 , $\epsilon_1 > \epsilon_2 > 0 \Rightarrow \bar{U}_{\epsilon_1}^\circ \subseteq U_{\epsilon_2}^\circ$.

Consider any class of open sets $\mathcal{U}_1 \subseteq \mathcal{U}(X)$.

Definition 3.2. We say that \mathcal{U}_1 is a tribe if and only if any open connected set which is a union of open sets in \mathcal{U}_1 , is again in \mathcal{U}_1 .

Clearly, both $\mathcal{U}(X)$ and $\mathcal{U}^r(X)$ are tribes, for all $r \geq 0$.

Assume now that \mathcal{U}_1 is endowed with some topology.

Definition 3.3. We say that \mathcal{U}_1 is a topological space of open sets if and only if

1. The topology of \mathcal{U}_1 is thinner than the Hausdorff distance topology, i.e. $\lim_{n \rightarrow \infty} U_n = U$ in $\mathcal{U}_1 \Rightarrow \lim_{n \rightarrow \infty} \rho_H(U_n, U) = 0$,
2. \mathcal{U}_1 admits outer approximations, in the sense that given $U \in \mathcal{U}_1$, there is an outer approximation $\{U_\epsilon^*\}_{\epsilon>0}$ of U in \mathcal{U}_1 such that $\lim_{\epsilon \rightarrow 0^+} U_\epsilon^* = U$.

We shall say that \mathcal{U}_1 admits inner approximations if and only if for every $U \in \mathcal{U}_1$, there is an inner approximation $\{U_\epsilon^\circ\}_\epsilon$ of U in \mathcal{U}_1 such that $\lim_{\epsilon \rightarrow 0^+} U_\epsilon^\circ = U$.

It follows, from item 1. of definition above, that the equivalence relation \sim is (topologically) closed. Therefore, if \mathcal{U}_1 is a topological space of open sets then the quotient $\mathbb{U}_1 = \mathcal{U}_1 / \sim$ is a Hausdorff topological space. The Hausdorff distance ρ_H is a pseudo-metric¹ on \mathcal{U}_1 , which induces a metric on \mathbb{U}_1 . Of course the quotient topology on \mathbb{U}_1 is thinner than the metric topology of (\mathbb{U}_1, ρ_H) .

¹We shall use the term pseudo-metric to any function $\rho : X \times X \rightarrow [0, +\infty]$ which is symmetric and satisfies the triangle inequality. The pseudo-metric ρ is called a metric if $\rho(x, y) = 0$ implies $x = y$.

We now topologize the sets $\mathcal{U}(X)$ and $\mathcal{U}^0(X)$, turning them into topological spaces of open sets.

Consider first the following pseudo-metric ρ on $\mathcal{U}(X)$. Given $U, V \in \mathcal{U}(X)$,

$$\rho(U, V) = \max \{ \rho_H(U, V), \rho_H(U^e, V^e) \} .$$

Clearly ρ satisfies $\rho_H(U, V) \leq \rho(U, V)$, and $\rho(U, V) = 0$ if and only if $U \sim V$.

Let us say that an open set $U \subseteq X$ has *regular boundary* if and only if $\partial U = \partial \bar{U}$. Notice that for each open set $U \in \mathcal{U}(X)$, \widehat{U} is the unique open set with regular boundary of its equivalence class.

Proposition 3.1. *Let $U \in \mathcal{U}(X)$.*

1. *Defining $U_\epsilon^* = B_\epsilon(U)$, the family $\{U_\epsilon^*\}_{\epsilon>0}$ is an outer approximation of U .*
2. *Defining U_ϵ° to be the largest connected component of $\{x \in X : d(x, U^c) > \epsilon\}$, this is an open set with regular boundary and $\{U_\epsilon^\circ\}_{\epsilon>0}$ is an inner approximation of U .*

Furthermore, for any inner approximation $\{U_\epsilon^\circ\}_{\epsilon>0}$, resp. outer approximation $\{U_\epsilon^\}_{\epsilon>0}$, of U in $\mathcal{U}(X)$, $\lim_{\epsilon \rightarrow 0^+} \rho(U_\epsilon^\circ, U) = 0 = \lim_{\epsilon \rightarrow 0^+} \rho(U_\epsilon^*, U)$. In particular, with the topology associated to ρ , $\mathcal{U}(X)$ is a topological space of open sets, which admits inner approximations.*

Next proposition characterizes the topology defined by ρ .

Proposition 3.2. *Given $U, V \in \mathcal{U}(X)$, and $\epsilon > 0$,*

$$\rho(U, V) \leq \epsilon \implies U_\epsilon^\circ \subseteq \widehat{V} \subseteq U_\epsilon^* .$$

Proof. First $\rho_H(U, V) \leq \epsilon$ implies that $\widehat{V} \subseteq \bar{V} \subseteq B_\epsilon(U) = U_\epsilon^*$. On the other hand $\rho_H(U^e, V^e) \leq \epsilon$ implies that $V^e \subseteq B_\epsilon(U^e) \subseteq B_\epsilon(U^c)$, which in turn implies that $U_\epsilon^\circ \subseteq \widehat{V}$. □

The quotient space $\mathbb{U}(X)$ can be identified with the subspace of $\mathcal{U}(X)$ formed by all open sets which have regular boundary. With this identification, metric ρ is given by

$$\rho(U, V) = \max \{ \rho_H(U, V), \rho_H(U^c, V^c) \} .$$

Notice that ρ may take infinite values. For any open set $U \in \mathcal{U}(X)$, $\rho(U, X) = \rho_H(U^c, \emptyset) = +\infty$.

Proposition 3.3. *With Hausdorff distance ρ_H , $\mathcal{U}^r(X)$, resp. $\mathbb{U}^r(X)$, is a compact pseudo-metric, resp. metric, space.*

Proof. Denote by $\mathbb{K}(X)$ the space of all non-empty compact subsets of X . With the Hausdorff distance ρ_H , $\mathbb{K}(X)$ is a compact metric space. See the Blaschke selection theorem in [F]. Therefore it is enough to check that the correspondence $U \mapsto \bar{U}$, from $\mathcal{U}^r(X)$ into $\mathbb{K}(X)$ induces a one-to-one mapping from $\mathbb{U}^r(X)$ onto a closed subspace of $\mathbb{K}(X)$. \square

We consider on $\mathcal{U}^0(X)$ the inductive limit topology of the compact pseudo-metric spaces $(\mathcal{U}^r(X), \rho_H)$ as $r \rightarrow 0^+$. Then the quotient topology on $\mathbb{U}^0(X)$ is again the inductive limit topology of the compact metric spaces $(\mathbb{U}^r(X), \rho_H)$.

Proposition 3.4. *With notation of proposition 3.1, for any $U \in \mathcal{U}^r(X)$ we have $U_\epsilon^* \in \mathcal{U}^r(X)$ and $U_\epsilon^c \in \mathcal{U}^{r-\epsilon}(X)$. For each $r \geq 0$, $\mathcal{U}^r(X)$ is a topological space of open sets. For $r = 0$, the inductive limit $\mathcal{U}^0(X) = \lim_{r \rightarrow 0^+} \mathcal{U}^r(X)$ also admits inner approximations.*

Notice that $\mathcal{U}^r(X)$ does not admit inner approximations for $r > 0$, since the open set $U = B_r(x) \in \mathcal{U}^r(X)$ fails to have inner approximations.

Next we shall characterize the compact topologies on $\mathcal{U}^r(X)$. We need the following fact:

Proposition 3.5. *There is some constant C_r , depending on $r > 0$ such that for all $U \in \mathcal{U}^r(X)$,*

$$H^{n-1}(\partial U) \leq C_r m(U) .$$

In particular, every open set $U \in \mathcal{U}^r(X)$ is Jordan measurable, i.e. $m(\partial U) = 0$.

Consider the pseudo-metrics ρ_H and ρ_m . It is obvious that $\rho_H(U, V) = 0 \Leftrightarrow \bar{U} = \bar{V}$. Analogously, using that both U and V are Jordan measurable open sets, one easily checks that $\rho_m(U, V) = 0 \Leftrightarrow \bar{U} = \bar{V}$. Therefore both ρ_H and ρ_m induce metrics on the quotient $\mathbb{U}^r(X)$.

Proposition 3.6. *For each $r > 0$ there is some constant C_r such that for any pair of open sets $U, V \in \mathcal{U}^r(X)$, $\rho_m(U, V) \leq C_r \rho_H(U, V)$. The two metrics induce the same compact topology on $\mathbb{U}^r(X)$.*

Proof. There is some constant K_r such that $H^{n-1}(\partial U) \leq K_r$ for all $U \in \mathcal{U}^r(X)$. Letting $\epsilon = \rho_H(U, V)$, $\rho_m(U, V)$ is bounded by the maximum volume of an ϵ -neighbourhood of both ∂U and ∂V , which is of order ϵK_r . It follows, using a standard abstract topology argument that the two metrics define the same Hausdorff compact topology on the quotient space. \square

Corollary 3.7. *The real valued function $U \mapsto m(U)$ is continuous over $\mathbb{U}^0(X)$.*

It is enough to prove proposition 3.5 when $X = \mathbb{R}^n$ and d is the Euclidean distance. In general we cover X with an atlas formed by a finite number of charts. Any geodesically r -regular open set can be decomposed as the union of a finite number of open sets in $\mathcal{U}^r(X)$, each of them completely covered by a single chart in the chosen atlas. Then there is some $0 < r' < r$ such that in each of these charts a geodesically r -regular open set is r' -regular w.r.t. the Euclidean metric.

Lemma 3.8. *In a plane convex trapezium with both diagonals of the same length $r > 0$ there is a pair of adjacent edges of length less the r .*

Proof. There is at least an obtuse angle in the trapezium. Compare the edges incident at this vertex with the opposed diagonal. \square

Given a subset $A \subseteq \{x \in \mathbb{R}^n : d(x, a) = r\}$ define the sector

$$S(a, A) = \{(1-t)a + tx : t \in [0, 1[, x \in A\}.$$

Lemma 3.9. *Let $\Omega = B_r(a) \cup B_r(b) \subseteq \mathbb{R}^n$, and consider two subsets $A, B \subseteq \partial\Omega$ with $A \subseteq \{x \in \mathbb{R}^n : d(x, a) = r\}$ and $B \subseteq \{x \in \mathbb{R}^n : d(x, b) = r\}$.*

If $A \cap B = \emptyset$ then $S(a, A) \cap S(b, B) = \emptyset$.

Proof. If $S(a, A) \cap S(b, B) \neq \emptyset$ then there are points $x \in A$ and $y \in B$ such that the straight segments $[a, x]$ and $[b, y]$ intersect at some point. Consider the plane trapezium with vertices a, y, x and b . This is a plane convex trapezium because the two diagonals $[a, x]$ and $[b, y]$ cross each other. By the previous lemma either $d(a, y) < r$ or $d(b, x) < r$ which in any case contradicts the fact that $x, y \in \partial\Omega$. \square

Lemma 3.10. For all $U \in \mathcal{U}^r(\mathbb{R}^n)$,

$$H^{n-1}(\partial U) \leq \frac{\sigma_n}{r} m(U) ,$$

where σ_n is the ratio between the $(n-1)$ -volume of $\{x \in \mathbb{R}^n : \|x\| = 1\}$ and the n -volume of $\{x \in \mathbb{R}^n : \|x\| \leq 1\}$.

Proof. Denote by H^{n-1} the $(n-1)$ -Hausdorff measure in \mathbb{R}^n , and by V^n the Lebesgue measure in \mathbb{R}^n . Then for any sector $S(a, A)$, with $A \subseteq \{x \in \mathbb{R}^n : d(x, a) = r\}$, we have

$$\frac{H^{n-1}(A)}{V^n(S(a, A))} = \frac{\sigma_n}{r} .$$

Claim: For any finite union $\Omega = \bigcup_{i=1}^m B_r(x_i)$,


$$H^{n-1}(\partial\Omega) \leq \frac{\sigma_n}{r} V^n(\Omega) .$$

We can cover $\partial\Omega$ with pairwise disjoint measurable sets A_i , each contained in the sphere $\{x \in \mathbb{R}^n : d(x, x_i) = r\}$. Then

$$\begin{aligned} H^{n-1}(\partial\Omega) &= \sum_{i=1}^m H^{n-1}(A_i) \\ &= \frac{\sigma_n}{r} \sum_{i=1}^m V^n(S(x_i, A_i)) \\ &\leq \frac{\sigma_n}{r} V^n(\Omega) , \end{aligned}$$

since the sectors $S(x_i, A_i) \subseteq \Omega$ are pairwise disjoint.

In general, if Ω is an infinite union of open balls of radius $r > 0$, the proof becomes more involved using Geometric Measure Theory techniques, and will be omitted here. Just to give an idea, notice that complements of r -regular open sets are compact sets, which are r -convex, in the sense that they are intersections of an arbitrary number of open ball complements $\{x \in \mathbb{R}^n : d(x, a) \geq r\}$. These r -convex sets share with convex sets some nice properties. Given any r -convex set $K \subseteq \mathbb{R}^n$ with connected complement there is $\delta_0(K) > 0$ and a Lipschitz projection $p : B_{\delta_0}(K) \rightarrow K$ such that $d(x, K) = \|x - p(x)\|$, for all $x \in B_{\delta_0}(K)$. Furthermore $\varphi(x) = d^2(x, K)$ is a class C^1 function with gradient $\nabla\varphi(x) = 2(x - p(x))$. These objects can be used to find Lipschitz parametrizations of the boundary $\partial\Omega$. \square



4 Topological Semigroups of Open Maps

Let $\mathcal{S}(X)$ be the space of all point-set maps on X , that is, $\varphi \in \mathcal{S}$ if and only if φ is a map from X into the power set of X , i.e. $\varphi : X \rightarrow \mathcal{P}(X)$. For a point-set map $\varphi \in \mathcal{S}$ and a subset $A \subseteq X$ the *image* $\varphi(A) \in \mathcal{P}(X)$ is defined by $\varphi(A) = \cup_{x \in A} \varphi(x)$. We say that $A \subseteq X$ is φ -*invariant* when $\varphi(A) \subseteq A$. Analogously, we say that A is *fully φ -invariant* if $\varphi(A) = A$. For two point-set maps $\varphi, \psi \in \mathcal{S}$ the usual composition product $\varphi \circ \psi : X \rightarrow \mathcal{P}(X)$ of φ and ψ at x is defined by

$$(\varphi \circ \psi)(x) = \varphi(\psi(x)) = \cup_{y \in \psi(x)} \varphi(y) .$$

Clearly, $\mathcal{S}(X)$ with this composition product is a semigroup.

We say that a point-set map $\varphi \in \mathcal{S}(X)$ is *open* when

$$\text{graph } \varphi = \{(x, y) \in X \times X : y \in \varphi(x)\}$$

is an open subset of $X \times X$.

We define $\mathcal{O}(X)$ to be the space of all point-set maps φ in $\mathcal{S}(X)$ that satisfy:

1. φ is open, and

2. $\varphi(x)$ is non-empty and connected for every $x \in X$,

that is, $\varphi \in \mathcal{O}(X)$ if and only if φ is an open point-set map on X with connected non-empty values.

Proposition 4.1. *Given $\varphi \in \mathcal{O}(X)$ and an open set $C \subseteq X$, if C is connected then $\varphi(C)$ is open and connected.*

Proof. Let $C \subseteq X$ be an open connected set. It is clear that $\varphi(C)$ is open, since φ is open. Assume, by contradiction, that $Y = \varphi(C)$ is not connected. There are then non-empty, disjoint, open sets A and B such that $Y = A \cup B$. Clearly,

$$\varphi^{-1}(A) = \{x \in C : \varphi(x) \cap A \neq \emptyset\} \quad \text{and} \quad \varphi^{-1}(B) = \{x \in C : \varphi(x) \cap B \neq \emptyset\}$$

are non-empty open subsets of C whose union is C . Let us show that $\varphi^{-1}(A)$ and $\varphi^{-1}(B)$ are disjoint. Again, by contradiction, we suppose there is $x \in C$ such that $\varphi(x) \cap A \neq \emptyset$ and $\varphi(x) \cap B \neq \emptyset$. Since A and B are disjoint, $\varphi(x)$ is not connected and we get a contradiction. Thus, $\varphi^{-1}(A)$ and $\varphi^{-1}(B)$ are disjoint. We conclude that C is not connected. We have reached a contradiction, so $\varphi(C)$ is connected. \square

From the previous proposition it follows easily that

Proposition 4.2. *$\mathcal{O}(X)$ is a subsemigroup of $\mathcal{S}(X)$.*

Given $\varphi \in \mathcal{O}(X)$ we define $\overline{\varphi}, \widehat{\varphi} : X \rightarrow \mathcal{P}(X)$ by setting

$$\text{graph } \overline{\varphi} = \overline{\text{graph } \varphi} \quad \text{and}$$

$$\text{graph } \widehat{\varphi} = (\overline{\text{graph } \varphi})^\circ .$$

We say that $\varphi_1, \varphi_2 \in \mathcal{O}(X)$ are equivalent, and write $\varphi_1 \sim \varphi_2$, if and only if $\overline{\varphi_1} = \overline{\varphi_2}$, or, which is equivalent, $\widehat{\varphi_1} = \widehat{\varphi_2}$.

Proposition 4.3. *For all $\varphi, \psi \in \mathcal{O}(X)$, $\varphi \circ \psi \sim \widehat{\varphi} \circ \widehat{\psi}$.*

Proof. It is enough to prove that $\widehat{\varphi} \circ \psi \sim \varphi \circ \psi \sim \varphi \circ \widehat{\psi}$. To prove the first equivalence note that $\text{graph}(\varphi \circ \psi) \subseteq \text{graph}(\widehat{\varphi} \circ \psi)$. Thus it is enough to see that first of these graphs is dense in the second. Take $(x, z) \in \text{graph}(\widehat{\varphi} \circ \psi)$. Then there is $y \in \psi(x)$ such that $z \in \widehat{\varphi}(y)$. Since $\text{graph}(\varphi)$ is dense in $\text{graph}(\widehat{\varphi})$, there is a sequence $(y_n, z_n) \in \text{graph}(\varphi)$ converging to (y, z) . Because $\psi(x)$ is open, we have $y_n \in \psi(x)$ for all large n , which proves that $(x, z_n) \in \text{graph}(\varphi \circ \psi)$ approximates (x, z) as n tends to infinity. Therefore $\text{graph}(\varphi \circ \psi)$ is dense in $\text{graph}(\widehat{\varphi} \circ \psi)$.

In a similar way, we can prove that $\text{graph}(\varphi \circ \psi)$ is dense in $\text{graph}(\varphi \circ \widehat{\psi})$, which implies the second equivalence. □

From this proposition follows that the relation \sim is compatible with composition. In particular the quotient $\mathbb{O}(X) = \mathcal{O}(X)/\sim$ inherits a semigroup structure from the usual composition. As a set, $\mathbb{O}(X)$ can be identified with the subspace of all $\varphi \in \mathcal{O}(X)$ such that $\text{graph}(\varphi)$ is an open set with regular boundary. We shall say that such open maps φ *have regular boundary*. Notice that compositions in $\mathbb{O}(X)$ and $\mathcal{O}(X)$ are different. Given $\varphi, \psi \in \mathbb{O}(X)$, regarded as open maps with regular boundary, if $\varphi \circ \psi = \zeta$ in $\mathcal{O}(X)$ then $\varphi \circ \psi = \widehat{\zeta}$ in $\mathbb{O}(X)$.

From definition 3.2, it is clear that

Proposition 4.4. *Given any tribe of open sets $\mathcal{U}_1 \subseteq \mathcal{U}(X)$ the set*

$$\mathcal{O}_1 = \{ \varphi \in \mathcal{O}(X) : \forall x \in X, \varphi(x) \in \mathcal{U}_1 \}$$

is a subsemigroup of $\mathcal{O}(X)$.

Given $r > 0$, we say that a map $\varphi \in \mathcal{O}(X)$ is r -regular if for every $x \in X$ the open set $\varphi(x)$ is r -regular, i.e. $\varphi(x) \in \mathcal{U}^r(X)$. We shall denote by $\mathcal{O}^r(X)$ the subset of all r -regular open maps. For $r = 0$ we set $\mathcal{O}^0(X) = \bigcup_{r>0} \mathcal{O}^r(X)$. Since each $\mathcal{O}^r(X)$ is a tribe, it follows from previous proposition that:

Proposition 4.5. *For all $r \geq 0$, $\mathcal{O}^r(X)$ is a subsemigroup of $\mathcal{O}(X)$*

Off course setting $\mathbb{O}^r(X) = \mathcal{O}^r(X)/\sim$, this is a subsemigroup of $\mathbb{O}(X)$.

We say that a point-set map is *Lipschitz* if and only if the map $x \mapsto \varphi(x)$ is Lipschitz with respect to the Hausdorff distance ρ_H , i.e. there is $K > 0$ such that

$\rho_H(\varphi(x), \varphi(y)) \leq K d(x, y)$ for every $y \in X$. We shall denote by $\text{Lip}(\varphi)$ the greatest lower bound of all Lipschitz constants K for this map. We denote by $\mathcal{O}_{\text{Lip}}(X)$ the subset of all Lipschitz point-set maps in $\mathcal{O}(X)$.

We recall the following continuity concepts. A point-set map $\varphi : X \rightarrow \mathcal{P}(X)$ with non-empty values is called *lower semi-continuous*, respectively *upper semi-continuous*, if for every $x \in X$ and $\epsilon > 0$ there is a neighborhood N_x of x in X such that for any $y \in N_x$, $\varphi(x) \subseteq B_\epsilon(\varphi(y))$, respectively $\varphi(y) \subseteq B_\epsilon(\varphi(x))$. It is called *continuous* if $\varphi : X \rightarrow \mathcal{P}(X)$ is both lower semi-continuous and upper semi-continuous. This means continuity with respect to the Hausdorff distance. We denote by $\mathcal{O}_{\text{Cont}}(X)$ the subset of all continuous point-set maps in $\mathcal{O}(X)$.

Proposition 4.6. $\mathcal{O}_{\text{Lip}}(X)$ and $\mathcal{O}_{\text{Cont}}(X)$ are both subsemigroups of $\mathcal{O}(X)$.

Proof. Let $\varphi, \psi \in \mathcal{O}_{\text{Lip}}$ with constants $K = \text{Lip}(\varphi)$ and $M = \text{Lip}(\psi)$. Take $x_1, x_2 \in X$ with $r = d(x_1, x_2)$. It is enough to prove that $(\psi \circ \varphi)(x_2) \subseteq \overline{B}_{MKr}((\psi \circ \varphi)(x_1))$. Then, by symmetry, exchanging the roles of x_1 and x_2 we get

$$\rho_H((\psi \circ \varphi)(x_1), (\psi \circ \varphi)(x_2)) \leq MKr = MKd(x_1, x_2).$$

Given $z_2 \in (\psi \circ \varphi)(x_2)$ there is $y_2 \in \varphi(x_2)$ such that $z_2 \in \psi(y_2)$. Since φ is Lipschitz, with $\text{Lip}(\varphi) = K$, we get $y_2 \in \overline{B}_{Kr}(\varphi(x_1))$. Therefore, there is $y_1 \in \varphi(x_1)$ such that $d(y_2, y_1) \leq Kr$. Because ψ is also Lipschitz, with $\text{Lip}(\psi) = M$, we get

$$z_2 \in \overline{B}_{MKr}(\psi(y_1)) \subseteq \overline{B}_{MKr}((\psi \circ \varphi)(x_1)).$$

□

We shall denote by $\mathbb{O}_{\text{Lip}}(X)$ and $\mathbb{O}_{\text{Cont}}(X)$ the corresponding subsemigroups of $\mathbb{O}(X)$. For any $r \geq 0$, define the semigroups $\mathcal{O}_{\text{Lip}}^r = \mathcal{O}_{\text{Lip}} \cap \mathcal{O}^r$ and $\mathcal{O}_{\text{Cont}}^r = \mathcal{O}_{\text{Cont}} \cap \mathcal{O}^r$. Accordingly, their quotients will be denoted by $\mathbb{O}_{\text{Lip}}^r$ and $\mathbb{O}_{\text{Cont}}^r$.

The following proposition is easily proved.

Proposition 4.7. 1. All maps φ in $\mathcal{O}(X)$ are lower semi-continuous.

2. For every map $\varphi \in \mathcal{O}(X)$, $\overline{\varphi}$ is upper semi-continuous.

3. Every map $\varphi \in \mathcal{O}_{\text{Lip}}(X)$ is continuous.

Identifying each $\varphi \in \mathcal{O}(X)$ with its graph $\text{graph}(\varphi)$ we can see $\mathcal{O}(X)$, respectively $\mathbb{O}(X)$, as a subset of $\mathcal{U}(X \times X)$, resp. $\mathbb{U}(X \times X)$. We are going to consider some topologies on the semigroups $\mathcal{O}(X)$, $\mathcal{O}_{\text{Lip}}^r(X)$, $\mathbb{O}(X)$ and $\mathbb{O}_{\text{Lip}}^r(X)$. Let us now give some abstract definitions:

Definition 4.1. *We say that a family of open point-set maps $\{\varphi_\epsilon\}_{\epsilon>0}$ is an outer approximation, respectively inner approximation, of $\varphi \in \mathcal{O}$ when $\{\text{graph} \varphi_\epsilon\}_{\epsilon>0}$ is an outer, respectively inner, approximation of $\text{graph} \varphi$.*

Given open maps $\varphi, \psi : X \rightarrow \mathcal{P}(X)$, we shall write $\varphi \leq \psi$ to mean that $\text{graph}(\varphi) \subseteq \text{graph}(\psi)$, and $\varphi \prec \psi$ to say that $\overline{\text{graph}(\varphi)} \subseteq \text{graph}(\psi)$. Clearly, these are transitive relations compatible with compositions.

Consider any subsemigroup of open maps $\mathcal{O}_1 \subseteq \mathcal{O}(X)$, endowed with some topology.

Definition 4.2. *We say that \mathcal{O}_1 is a topological semigroup of open maps if and only if*

- (a) \mathcal{O}_1 is a topological space of open sets in $\mathcal{U}(X \times X)$, in the sense of definition 3.3.
- (b) Given $\epsilon > 0$, an integer $N \in \mathbb{N}$, and non-empty open subsets $U, V \subseteq X$ such that $\overline{U \times V} \subseteq \text{graph} \varphi^N$, there is a neighbourhood \mathcal{N} of φ in \mathcal{O}_1 such that for all $\psi \in \mathcal{N}$ and $x \in \overline{U}$, $m(V \setminus \widehat{\psi^N}(x)) < \epsilon$.

If \mathcal{O}_1 is a topological semigroup of open maps, we shall say that the quotient semigroup $\mathbb{O}_1 = \mathcal{O}_1 / \sim$ is also a topological semigroup of open maps. Alternatively, regarding elements in \mathbb{O}_1 as open maps with regular boundary, a similar definition as above can be used to characterize when is \mathbb{O}_1 a topological semigroup of open maps. Notice that condition (b) becomes even simpler because, in the quotient semigroup, ψ^N will play the role that $\widehat{\psi^N}$ has in \mathcal{O}_1 .

We say that a topological semigroup \mathcal{O}_1 is lower semi-continuous if and only if given inner approximations in \mathcal{O}_1 , $\{\varphi_\epsilon^\circ\}_{\epsilon>0}$ and $\{\psi_\epsilon^\circ\}_{\epsilon>0}$ of $\varphi, \psi \in \mathcal{O}_1$, respectively, if $\lim_{\epsilon \rightarrow 0^+} \varphi_\epsilon^\circ = \varphi$ and $\lim_{\epsilon \rightarrow 0^+} \psi_\epsilon^\circ = \psi$, then $\lim_{\epsilon \rightarrow 0^+} \psi_\epsilon^\circ \circ \varphi_\epsilon^\circ = \psi \circ \varphi$.

When the composition mapping is continuous we shall say that \mathcal{O}_1 is continuous.

Proposition 4.8. *Given a topological semigroup of open maps \mathbb{O}_1 and $N \in \mathbb{N}$, the family of mappings $\varphi \mapsto \varphi^N(x)$ from \mathbb{O}_1 into $(\mathbb{U}(X), \rho_H)$ is lower semi-equicontinuous, i.e. given $\epsilon > 0$ there is a neighbourhood \mathcal{N} of φ in \mathbb{O}_1 such that for all $\psi \in \mathcal{N}$ and $x \in X$, $B_\epsilon(\psi^N(x)) \supseteq \varphi^N(x)$.*

Proof. Given $\epsilon > 0$ cover $\overline{\text{graph } \varphi^N}$ by a finite number of ϵ -radius balls $B_\epsilon(x_i, y_i) = B_\epsilon(x_i) \times B_\epsilon(y_i)$, $i = 1, \dots, m$. Then for each i , choose a neighbourhood \mathcal{N}_i of φ in \mathbb{O}_1 such that for all $\psi \in \mathcal{N}_i$ and $x \in \overline{B_\epsilon(x_i)}$, $m(B_\epsilon(y_i) \setminus \psi^N(x)) < m(B_\epsilon(y_i))$, which in turn implies that $\psi^N(x) \cap B_\epsilon(y_i) \neq \emptyset$. Given $\psi \in \cap_{i=1}^m \mathcal{N}_i$, $x \in X$ and $y \in \varphi^N(x)$, one of the points (x_i, y_i) is such that $(x, y) \in B_\epsilon(x_i, y_i)$. Taking $z \in \psi^N(x) \cap B_\epsilon(y_i)$ we see that $d(y, z) \leq d(y, y_i) + d(y_i, z) \leq 2\epsilon$, which implies $y \in B_{2\epsilon}(\psi^N(x))$. This proves that $\varphi^N(x) \subseteq B_{2\epsilon}(\psi^N(x))$. \square

Proposition 4.9. *Given $\varphi \in \mathcal{O}(X)$, define φ_ϵ^* by*

$$\text{graph } (\varphi_\epsilon^*) = B_\epsilon(\text{graph } (\varphi)) ,$$

and define φ_ϵ° setting $\varphi_\epsilon^\circ(x)$ to be the largest connected component of the open set

$$\{ y \in X : d((x, y), \text{graph } (\varphi)^c) > \epsilon \} .$$

Then $\varphi_\epsilon^ \in \mathcal{O}(X)$, and $\varphi_\epsilon^\circ \in \mathcal{O}(X)$ for all small enough $\epsilon > 0$. Moreover $\{\varphi_\epsilon^\circ\}_{\epsilon>0}$ and $\{\varphi_\epsilon^*\}_{\epsilon>0}$ are respectively inner and outer approximations of φ in $\mathcal{O}(X)$.*

It should be clear by definition that compositions of inner approximations is still an inner approximation.

Lemma 4.10. *Given $\varphi, \psi \in \mathcal{O}(X)$, if $\{\varphi_\epsilon^\circ\}_{\epsilon>0}$ and $\{\psi_\epsilon^\circ\}_{\epsilon>0}$ are inner approximations in $\mathcal{O}(X)$ of φ and ψ , respectively, then $\{\psi_\epsilon^\circ \circ \varphi_\epsilon^\circ\}_{\epsilon>0}$ is an inner approximation of $\psi \circ \varphi$.*

Given $\varphi \in \mathcal{O}$ and $\epsilon > 0$, we shall call ϵ -implosion, resp. ϵ -explosion to the inner and outer approximations defined in the previous proposition.

We now topologize the semigroups $\mathcal{O}(X)$ and $\mathcal{O}_{\text{Lip}}^0(X)$, turning them into topological semigroups of open maps. First consider $\mathcal{O}(X)$ as a subset of $\mathcal{U}(X \times X)$ and let ρ be the induced pseudo-metric, which is given by $\rho(\varphi, \psi) = \rho(\text{graph } (\varphi), \text{graph } (\psi))$.

The following characterization of the ϵ -ball for the pseudo-metric ρ follows from 3.2.

Proposition 4.11. *Given $\epsilon > 0$, for every $\varphi, \psi \in \mathcal{O}(X)$,*

$$\rho(\varphi, \psi) < \epsilon \implies \psi_\epsilon^\circ \prec \widehat{\varphi} \prec \psi_\epsilon^*.$$

Proposition 4.12. *With the topology associated to ρ , $\mathcal{O}(X)$ is a topological semigroup of open maps, which admits inner approximations and is lower semi-continuous.*

Proof. Item (a) of definition 4.2 follows from proposition 3.1. By proposition 4.9, $\mathcal{O}(X)$ admits inner approximations. From lemma 4.10 and proposition 3.1 we see that $\mathcal{O}(X)$ is lower semi-continuous. To see item (b), let $U, V \subseteq X$ be non-empty open sets such that $\overline{U \times V} \subseteq \text{graph } \varphi^N$. Taking $\delta > 0$ small enough we have $\overline{U \times V} \subseteq \text{graph } (\varphi_\delta^\circ)^N$. Consider the δ -neighbourhood $\mathcal{N} = B_\delta(\varphi)$ w.r.t. the pseudo-metric ρ . If $\psi \in \mathcal{N}$ then, by proposition 4.11, $\varphi_\delta^\circ \prec \widehat{\psi}$, implying that

$$\overline{U \times V} \subseteq \text{graph } (\varphi_\delta^\circ)^N \subseteq \text{graph } (\widehat{\psi})^N \subseteq \text{graph } \widehat{\psi}^N.$$

Therefore $m(V \setminus \widehat{\psi}^N(x)) = 0$ for all $x \in \overline{U}$. □

It is not difficult to see that composition is not continuous. In particular $\mathcal{O}(X)$ is not a continuous topological semigroup of open maps. We have however

Proposition 4.13. *With the topology associated to ρ , $\mathcal{O}_{\text{Cont}}(X)$ is a continuous topological semigroup of open maps.*

This proposition follows easily from the next lemma.

Lemma 4.14. *Given $\varphi \in \mathcal{O}_{\text{Cont}}(X)$ we have, using the notation of proposition 4.9, $\varphi_\epsilon^* \in \mathcal{O}_{\text{Cont}}(X)$, and $\varphi_\epsilon^\circ \in \mathcal{O}_{\text{Cont}}(X)$ for every small enough $\epsilon > 0$. Moreover, given $\varphi, \psi \in \mathcal{O}_{\text{Cont}}(X)$, if $\{\varphi_\epsilon^*\}_{\epsilon > 0}$ and $\{\psi_\epsilon^*\}_{\epsilon > 0}$ are outer approximations in $\mathcal{O}_{\text{Cont}}(X)$ of φ and ψ , respectively, then $\{\psi_\epsilon^* \circ \varphi_\epsilon^*\}_{\epsilon > 0}$ is an outer approximation of $\psi \circ \varphi$.*

In the sequell we introduce a semigroup topology on $\mathcal{O}_{\text{Lip}}^0(X)$. Consider in $\mathcal{O}(X) \subseteq \mathcal{U}(X \times X)$ the pseudo-metrics ρ_H, ρ_m, d_H and d_m , where the last two are defined by

Definition 4.3.

$$d_H(\varphi, \psi) = \sup_{x \in X} \rho_H(\varphi(x), \psi(x)) \quad \text{and}$$

$$d_m(\varphi, \psi) = \sup_{x \in X} \rho_m(\varphi(x), \psi(x)) .$$

Given $r > 0$ and $K < +\infty$, define

$$\mathcal{O}_K^r = \{ \varphi \in \mathcal{O}^r(X) : \text{Lip}(\varphi) \leq K \} .$$

Proposition 4.15. *All four metrics ρ_H , ρ_m , d_H and d_m induce the same compact topology on \mathcal{O}_K^r .*

Proof. In general one has $\rho_H(\varphi, \psi) \leq d_H(\varphi, \psi)$ over $\mathcal{O}(X)$. If φ and ψ are Lipschitz with Lipschitz constant K , then we can also prove that $d_H(\varphi, \psi) \leq K \rho_H(\varphi, \psi)$. Therefore, ρ_H and d_H are Lipschitz equivalent over \mathcal{O}_K^r .

It is clear that

$$\rho_m(\varphi, \psi) = \int_X \rho_m(\varphi(x), \psi(x)) dm(x) \leq d_m(\varphi, \psi) .$$

Using proposition 3.6, we see that $d_m \leq C_r d_H$ over \mathcal{O}^r for some constant C_r depending on $r > 0$.

It is now enough to prove the compactity of \mathcal{O}_K^r w.r.t. d_H . This follows by an argument as in Ascoli-Arzelá's theorem. \mathcal{O}_K^r can be seen as a subset of the compact product space $\mathcal{U}^r(X)^X$. Of course \mathcal{O}_K^r is closed and, therefore, compact for the product topology. The set \mathcal{O}_K^r is equicontinuous because all maps in it are Lipschitz with the same Lipschitz constant. Thus the product topology (pointwise convergence) coincides with the topology of d_H (uniform convergence), which proves the compactity of \mathcal{O}_K^r w.r.t. d_H . \square

We shall consider on $\mathcal{O}_{\text{Lip}}^0 = \bigcup_{r>0, K<\infty} \mathcal{O}_K^r$, respectively on $\mathcal{O}_{\text{Lip}}^r = \bigcup_{K<\infty} \mathcal{O}_K^r$ if $r > 0$, the inductive limit topology of the compact subspaces \mathcal{O}_K^r .

Proposition 4.16. *With the inductive limit topology above, for each $r \geq 0$, $\mathcal{O}_{\text{Lip}}^r$ is a continuous topological semigroup of open maps.*

Proof. Item (a) is clear. The continuity of composition follows from proposition 4.17 bellow. To see item (b), take $\epsilon > 0$ and let $U, V \subseteq X$ be non-empty open sets such that $\overline{U \times V} \subseteq \text{graph } \varphi^N$. Consider the neighbourhood $\mathcal{N} = \{ \psi : d_m(\psi^N, \varphi^N) < \epsilon \}$ of φ in $\mathcal{O}_{\text{Lip}}^r$. Then if $\psi \in \mathcal{N}$, for all $x \in \overline{U}$,

$$m(V \setminus \widehat{\psi^N}(x)) \leq m(V \setminus \psi^N(x)) \leq m(\varphi^N(x) \setminus \psi^N(x)) \leq d_m(\psi^N, \varphi^N) < \epsilon. \quad \square$$

Proposition 4.17. *The composition map $(\varphi, \psi) \mapsto \varphi \circ \psi$ from $\mathcal{O}_K^r \times \mathcal{O}_K^r$ into $\mathcal{O}_{K^2}^r$ is continuous w.r.t. the compact topologies above.*

Proof. Using the lemma bellow we see that

$$\begin{aligned} \rho_H(\varphi_1(\psi_1(x)), \varphi_2(\psi_2(x))) &\leq \rho_H(\varphi_1(\psi_1(x)), \varphi_1(\psi_2(x))) + \rho_H(\varphi_1(\psi_2(x)), \varphi_2(\psi_2(x))) \\ &\leq K \rho_H(\psi_1(x), \psi_2(x)) + d_H(\varphi_1, \varphi_2) \\ &\leq K d_H(\psi_1, \psi_2) + d_H(\varphi_1, \varphi_2). \end{aligned}$$

Therefore

$$d_H(\varphi_1 \circ \psi_1, \varphi_2 \circ \psi_2) \leq K d_H(\psi_1, \psi_2) + d_H(\varphi_1, \varphi_2),$$

which proves continuity of the composition. \square

Lemma 4.18. *For all open sets $U, V \subseteq X$, and all maps $\varphi, \psi \in \mathcal{O}(X)$:*

1. $\rho_H(\varphi(U), \varphi(V)) \leq \text{Lip}(\varphi) \rho_H(U, V)$
2. $\rho_H(\varphi(U), \psi(U)) \leq d_H(\varphi, \psi)$

5 Dynamics of Open Maps

Given $\varphi \in \mathcal{O}(X)$, any sequence $x_0, x_1, \dots, x_n \in X$ such that $x_i \in \varphi(x_{i-1})$ for $i = 1, \dots, n$ will be called an *orbit* of φ , and we shall say that x_n is an *iterate* of the state x_0 . Given $x, y \in X$ let us write $x \rightsquigarrow_\varphi y$ when there exists some $n \in \mathbb{N}$ such that $y \in \varphi^n(x)$. The relation \rightsquigarrow_φ is transitive. Moreover, \rightsquigarrow_φ is an open relation, since φ

is an open point-set map. Given $A, B \subseteq X$ we shall write $A \rightsquigarrow_\varphi B$ when $x \rightsquigarrow_\varphi y$ for some $x \in A$ and some $y \in B$. We define the *recurrent set* of φ , and denote it by $\Omega(\varphi)$, as the set of all states $x \in X$ such that x is an iterate of x , that is,

$$\Omega(\varphi) = \{x \in X : x \rightsquigarrow_\varphi x\}.$$

Clearly, $\Omega(\varphi)$ is an open set. On $\Omega(\varphi)$ the relation \rightsquigarrow_φ is reflexive and transitive and thus is a preorder. We call the elements of $\Omega(\varphi)$ *recurrent states*. The recurrent set $\Omega(\varphi)$ can be decomposed into classes defined by the following equivalence relation. Given $x, y \in X$ we shall write $x \rightsquigarrow_\varphi y$ when $x = y$, or then $x \rightsquigarrow_\varphi y$ and $y \rightsquigarrow_\varphi x$. Since \rightsquigarrow_φ is a preorder on $\Omega(\varphi)$, \rightsquigarrow_φ is an equivalence relation. The equivalence classes of recurrent states will be called Ω -*classes*. Every Ω -class is open, since \rightsquigarrow_φ is an open relation. We shall denote by $\Lambda^\Omega(\varphi)$ the set of all Ω -classes of φ .

The relation \rightsquigarrow_φ imposes a partial ordering on $\Lambda^\Omega(\varphi)$. Indeed the relation \rightsquigarrow_φ is a preorder on $\Lambda^\Omega(\varphi)$ and in addition for $A, B \in \Lambda^\Omega(\varphi)$, $A \rightsquigarrow_\varphi B$ and $B \rightsquigarrow_\varphi A$ together imply $A = B$. The maximal elements in $\Lambda^\Omega(\varphi)$ under \rightsquigarrow_φ will be called Ω -*final classes*. In other words $D \in \Lambda^\Omega(\varphi)$ is an Ω -final class if for all $C \in \Lambda^\Omega(\varphi)$

$$D \rightsquigarrow_\varphi C \Rightarrow C = D.$$

We will denote by $\Lambda_{\text{final}}^\Omega(\varphi)$ the set of all Ω -final components of φ .

A recurrent state x will be called a *final recurrent state* if every iterate of x still has some iterate which comes back to x , that is,

$$\Omega_{\text{final}}(\varphi) = \{x \in \Omega(\varphi) : \text{for every } y \in X, x \rightsquigarrow_\varphi y \Rightarrow y \rightsquigarrow_\varphi x\},$$

where we denote by $\Omega_{\text{final}}(\varphi)$ the set of all final recurrent states. We shall call *final recurrent set* to $\Omega_{\text{final}}(\varphi)$. Indeed this is the set where all the dynamics will eventually end up. It should be regarded as the union of all "attractors" of φ . Clearly, final recurrent states and Ω -final classes are related in the following obvious way:

Proposition 5.1. $\Omega_{\text{final}}(\varphi)$ is the union of all Ω -final classes in $C \in \Lambda_{\text{final}}^\Omega(\varphi)$, each Ω -final class being fully φ -invariant, i.e. $\varphi(C) = C$.

For a subset $C \subseteq X$ we define the following open sets:

$$W_\Omega^s(C, \varphi) = W_\Omega^s(C) = \{z \in X : z \rightsquigarrow_\varphi C\}, \text{ and}$$

$$W_{\Omega}^u(C, \varphi) = W_{\Omega}^u(C) = \{z \in X : C \rightsquigarrow_{\varphi} z\},$$

that we shall call, respectively, the Ω -stable and Ω -unstable sets of C .

The following proposition implies a uniform spread on images of open maps. It follows by compactness of X using a standard type of argument.

Proposition 5.2. *Given $\varphi \in \mathcal{O}(X)$ there is a map $F : X \rightarrow X$ and $\xi_0 > 0$ such that*

$$B_{\xi_0}(\text{graph}(F)) \subseteq \text{graph}(\varphi).$$

Proposition 5.3. *The partial ordered set $\Lambda_{\text{final}}^{\Omega}(\varphi)$ is finite, and every state $x \in X$ has some iterate in $\Omega_{\text{final}}(\varphi)$, i.e. $X = W_{\Omega}^s(\Omega_{\text{final}}(\varphi))$.*

In other words, $\Omega_{\text{final}}(\varphi)$ splits into a finite number of attractors, attracting everything.

Proof. By proposition 5.2 there is some $\xi_0 > 0$ such that any Ω -final class, being fully φ -invariant, contains a ball of radius $\xi_0 > 0$. Therefore the volumes of Ω -final classes are uniformly bounded from zero. Since X is compact it follows that $\Lambda_{\text{final}}^{\Omega}(\varphi)$ must be finite. The second statement of this proposition is a consequence of the following two lemmas. \square

Lemma 5.4. $X = W_{\Omega}^s(\Omega(\varphi))$.

Proof. Let F and $\xi_0 > 0$ be as given by proposition 5.2. Take $x_0 \in X$ and define $x_n = F^n(x_0)$ for every $n \in \mathbb{N}$. Let z be a sublimit of x_n . If $d(x_n, z) < \xi_0$ and $d(x_{n+p}, z) < \xi_0$, then $x_{n+1} \in \varphi(z)$ and $z \in \varphi(x_{n+p-1})$. So $z \rightsquigarrow_{\varphi} z$ and thus $z \in \Omega(\varphi)$. Since $x_0 \rightsquigarrow_{\varphi} x_{n+p-1}$ and $x_{n+p-1} \rightsquigarrow_{\varphi} z$, we have $x_0 \rightsquigarrow_{\varphi} z$. \square

Lemma 5.5. *For each Ω -class $D \in \Lambda^{\Omega}(\varphi)$ there is some Ω -final class $D_0 \in \Lambda_{\text{final}}^{\Omega}(\varphi)$ such that $D \rightsquigarrow_{\varphi} D_0$.*

Proof. By Zorn's Lemma it is enough to prove that each forward chain $\mathcal{D} \subseteq \Lambda^{\Omega}(\varphi)$ has some upper bound $C \in \Lambda^{\Omega}(\varphi)$, i.e. for all $D \in \mathcal{D}$, $D \rightsquigarrow_{\varphi} C$. A forward chain \mathcal{D} in

$(\Lambda^\Omega(\varphi), \rightsquigarrow_\varphi)$ is a subset $\mathcal{D} \subseteq \Lambda^\Omega(\varphi)$ which is well ordered ² by \rightsquigarrow_φ . Assume there is no $z \in X$ such that for all $D \in \mathcal{D}$, $D \rightsquigarrow_\varphi z$. Define for each $D \in \mathcal{D}$,

$$S_D = \bigcap_{C \in \mathcal{D}, C \rightsquigarrow_\varphi D} W_\Omega^u(C) \setminus W_\Omega^u(D).$$

Then the family $\{S_D\}_{D \in \mathcal{D}}$ is formed of pairwise disjoint sets, and so we have $\sum_{D \in \mathcal{D}} m(S_D) < \infty$. Thus, there is some $D \in \mathcal{D}$ such that

$$m \left(\bigcup_{C \not\rightsquigarrow_\varphi D} S_C \right) = \sum_{C \not\rightsquigarrow_\varphi D} m(S_C) < c_0,$$

where c_0 is the m -volume of a ball of radius ξ_0 , for ξ_0 as in proposition 5.2. By the above assumption, we have $\bigcup_{C \not\rightsquigarrow_\varphi D} S_C = \bigcap_{C \rightsquigarrow_\varphi D} W_\Omega^u(C)$. Clearly this set is φ -invariant which contradicts the spread property of φ given by proposition 5.2. Therefore our assumption must be false and so there is some $z \in X$ such that for all $D \in \mathcal{D}$, $D \rightsquigarrow_\varphi z$. By proposition 5.4 there is some class $C \in \Lambda^\Omega(\varphi)$ such that $z \rightsquigarrow_\varphi C$. Thus, for all $D \in \mathcal{D}$, $D \rightsquigarrow_\varphi C$. \square

A connected component of $\Omega(\varphi)$, respectively $\Omega_{\text{final}}(\varphi)$, will be called an Ω -*component*, respectively an Ω -*final component*, of φ . We denote respectively by $\Sigma^\Omega(\varphi)$ and $\Sigma_{\text{final}}^\Omega(\varphi)$ the sets of all Ω -components, and of all Ω -final components, of φ .

Theorem 5.1. *The set of Ω -final components $\Sigma_{\text{final}}^\Omega(\varphi)$ is finite. For every component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$, $\varphi(C) \in \Sigma_{\text{final}}^\Omega(\varphi)$. The mapping $\pi_\varphi : \Sigma_{\text{final}}^\Omega(\varphi) \rightarrow \Sigma_{\text{final}}^\Omega(\varphi)$, defined by $\pi_\varphi(C) = \varphi(C)$, is a bijection, which permutes cyclically the components of each Ω -final class.*

Proof. By proposition 5.1, every Ω -final class C satisfies $\varphi(C) = C$. Thus, for every component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$, since $\varphi(C)$ is also connected, $\varphi(C)$ must be another final component in $\Sigma_{\text{final}}^\Omega(\varphi)$. It follows by proposition 5.2, that every Ω -final component must contain some ball of minimum radius ξ_0 . Therefore $\Sigma_{\text{final}}^\Omega(\varphi)$ is finite. Because each component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$ is formed by recurrent states, there is some integer $n \in \mathbb{N}$ such that $\varphi^n(C) = C$. By definition of an Ω -class it follows that $\bigcup_{i=0}^{n-1} \varphi^i(C)$ is precisely the class of C . \square

²A well ordered set is a totally ordered set such that every non-empty subset has a first element.

We shall call *period* of an Ω -final class C to the number d of its connected components. We will refer to the period of a component as the period of its class.

Next we introduce chain recurrency and relate it with the former recurrency concept. Given $\varphi \in \mathcal{O}(X)$ and $\epsilon > 0$, φ_ϵ^* will denote the ϵ -explosion of φ as defined in the previous section. Any sequence $x_0, x_1, \dots, x_n \in X$ such that $x_i \in \varphi_\epsilon^*(x_{i-1})$ for $i = 1, \dots, n$ will be called an ϵ -pseudo orbit of φ , and we shall say that x_n is an ϵ -pseudo iterate of the state x_0 . If y is an ϵ -pseudo iterate of x for every $\epsilon > 0$, y is said a *pseudo iterate* of x . Given $x, y \in X$ let us write $x \rightarrow_\varphi y$ when $x \rightsquigarrow_{\varphi_\epsilon^*} y$ for every $\epsilon > 0$. Clearly the relation \rightarrow_φ is a transitive, closed relation. Given $A, B \subseteq X$ we shall write $A \rightarrow_\varphi B$ when $x \rightarrow_\varphi y$ for some $x \in A$ and some $y \in B$.

We define the *chain recurrent set* of φ , and denote it by $R(\varphi)$, as the set of all states $x \in X$ such that x is a pseudo iterate of x , that is,

$$R(\varphi) = \{ x \in X : x \rightarrow_\varphi x \} .$$

The set $R(\varphi) = \bigcap_{\epsilon > 0} \overline{\Omega(\varphi_\epsilon^*)}$ is a closed set in X . The closed relation \rightarrow_φ is a preorder on $R(\varphi)$. We call the elements of $R(\varphi)$ *chain recurrent states*.

The recurrent set $R(\varphi)$ can be decomposed into classes defined by the following equivalence relation. Given $x, y \in X$ we shall write $x \rightleftharpoons_\varphi y$ when $x = y$, or then $x \rightarrow_\varphi y$ and $y \rightarrow_\varphi x$. Since \rightarrow_φ is a preorder on $R(\varphi)$, $\rightleftharpoons_\varphi$ is an equivalence relation. The equivalence classes of recurrent states will be called *R-classes*. Every *R-class* is closed, since $\rightleftharpoons_\varphi$ is a closed relation. We shall denote by $\Lambda^R(\varphi)$ the set of all *R-classes* of φ .

The relation \rightarrow_φ imposes a partial ordering on $\Lambda^R(\varphi)$. Indeed the relation \rightarrow_φ is a preorder on $\Lambda^R(\varphi)$ and in addition for $A, B \in \Lambda^R(\varphi)$, if $A \rightarrow_\varphi B$ and $B \rightarrow_\varphi A$ then $A = B$. The maximal elements in $\Lambda^R(\varphi)$ under \rightarrow_φ will be called *R-final classes*. In other words $D \in \Lambda^R(\varphi)$ is an *R-final class* if for all $C \in \Lambda^R(\varphi)$

$$D \rightarrow_\varphi C \Rightarrow C = D.$$

We will denote by $\Lambda_{\text{final}}^R(\varphi)$ the set of all *R-final components* of φ .

A chain recurrent state x will be called a *final chain recurrent state* if every pseudo iterate of x still has some pseudo iterate which comes back to x , that is,

$$R_{\text{final}}(\varphi) = \{ x \in R(\varphi) : \text{for every } y \in X, x \rightarrow_{\varphi} y \Rightarrow y \rightarrow_{\varphi} x \},$$

where we denote by $R_{\text{final}}(\varphi)$ the set of all final chain recurrent states. We shall call *final chain recurrent set* to $R_{\text{final}}(\varphi)$. Clearly, final recurrent states and R -final classes are related in the following obvious way:

Proposition 5.6. *$R_{\text{final}}(\varphi)$ is the union of all R -final classes $C \in \Lambda_{\text{final}}^R(\varphi)$, and each such class is fully $\bar{\varphi}$ -invariant, i.e. $\bar{\varphi}(C) = C$.*

Proof. Given $C \in \Lambda_{\text{final}}^R(\varphi)$, clearly $\bar{\varphi}(C) \subseteq C$. Now, let $y \in C$, and assume, by contradiction, that $y \notin \bar{\varphi}(C)$. Then

$$C \times \{y\} \cap \text{graph} \bar{\varphi} = \emptyset.$$

Take $x_0 \in C$. For every $\epsilon > 0$, $C_{\epsilon} = \bigcup_{n=1}^{\infty} (\varphi_{\epsilon}^*)^n(x_0)$ is the Ω -class of φ_{ϵ}^* containing x_0 . Off course $C = \bigcap_{\epsilon > 0} C_{\epsilon}$ and $\{C_{\epsilon}\}_{\epsilon > 0}$ is an outer approximation of C . By compactness of $C \times \{y\}$ and $\text{graph} \bar{\varphi}$ there is some small enough $\epsilon > 0$ such that

$$C_{\epsilon} \times \{y\} \cap \text{graph} \varphi_{\epsilon}^* = \emptyset,$$

which implies that $y \notin \varphi_{\epsilon}^*(C_{\epsilon}) = C_{\epsilon}$. But this is impossible because $y \in C \subseteq C_{\epsilon}$. \square

Proposition 5.7. *Every R -final class $A \in \Lambda_{\text{final}}^R(\varphi)$ contains some Ω -final class $C \in \Lambda_{\text{final}}^{\Omega}(\varphi)$, $A \supseteq C$.*

Proof. Given $A \in \Lambda_{\text{final}}^R(\varphi)$, by proposition 5.3 there is some class $C \in \Lambda_{\text{final}}^{\Omega}(\varphi)$ such that $A \rightsquigarrow_{\varphi} C$, which in turn implies $A \rightarrow_{\varphi} C$. Thus $A \supseteq C$, since A is an R -final class. \square

A connected component of $R(\varphi)$, respectively $R_{\text{final}}(\varphi)$, will be called an *R -component*, respectively an *R -final component*, of φ . We denote respectively by $\Sigma^R(\varphi)$ and $\Sigma_{\text{final}}^R(\varphi)$ the sets of all R -components, and of all R -final components, of φ .

Theorem 5.2. *The set of R -final components $\Sigma_{\text{final}}^R(\varphi)$ is finite. For every component $A \in \Sigma_{\text{final}}^R(\varphi)$, $\bar{\varphi}(A) \in \Sigma_{\text{final}}^{\Omega}(\varphi)$. The mapping $\bar{\pi}_{\varphi} : \Sigma_{\text{final}}^R(\varphi) \rightarrow \Sigma_{\text{final}}^R(\varphi)$, defined by $\bar{\pi}_{\varphi}(A) = \bar{\varphi}(A)$, is a bijection, which permutes cyclically the components of each R -final class.*

Proof. Given any component $A \in \Sigma_{\text{final}}^R(\varphi)$, since $\bar{\varphi}(A)$ is connected, it is completely contained inside a single component $A' \in \Sigma_{\text{final}}^R(\varphi)$. Let us define $\bar{\pi}_\varphi : \Sigma_{\text{final}}^R(\varphi) \rightarrow \Sigma_{\text{final}}^R(\varphi)$, setting $\bar{\pi}_\varphi(A) = A'$. By proposition 5.1, $\bar{\pi}_\varphi$ is surjective. Therefore each component of $\Sigma_{\text{final}}^R(\varphi)$ contains some image $\bar{\varphi}(x)$ and, by proposition 5.2, must have a minimum volume. By compactness of X the set of all components $\Sigma_{\text{final}}^R(\varphi)$ is finite. Thus, $\bar{\pi}_\varphi$ must be one-to-one. Again by the same proposition 5.1, $\bar{\pi}_\varphi$ permutes the components of each R -final class. Finally, this implies that $\bar{\varphi}(A)$ is a whole component, for each $A \in \Sigma_{\text{final}}^R(\varphi)$. \square

We shall call *period* of an R -final class A to the number d of its connected components. We will refer to the period of a component as the period of its class.

For a subset $C \subseteq X$ we define the following closed sets:

$$W_R^s(C, \varphi) = W_R^s(C) = \{z \in X : z \rightarrow_\varphi C\}, \text{ and}$$

$$W_R^u(C, \varphi) = W_R^u(C) = \{z \in X : C \rightarrow_\varphi z\},$$

that we shall call, respectively, the R -stable and R -unstable sets of C . Next proposition shows that final classes behave as 'attractors' do.

Proposition 5.8. *Given a final class $C \in \Lambda_{\text{final}}^R(\varphi)$ and a φ -invariant open set $U \subseteq X$ such that $\bar{U} \cap R(\varphi) = C$, $\bigcap_{n=1}^{\infty} \varphi^n(U) \subseteq C$.*

The proposition is corollary of

Lemma 5.9. *Given a φ -invariant open set $U \subseteq X$,*

$$\bigcap_{n=1}^{\infty} \varphi^n(U) \subseteq W_R^u(\bar{U} \cap R(\varphi)).$$

Proof. Given $x \in \bigcap_{n=1}^{\infty} \varphi^n(U)$, pick for each $n \in \mathbb{N}$, $z_n \in U$ such that $x \in \varphi^n(z_n)$. Taking z to be a sublimit of $\{z_n\}$, we have $z \rightarrow_\varphi x$. Consider the compact set $\Gamma = W_R^u(z) \cap W_R^s(x) \subseteq \bar{U}$. This set is weakly invariant in the sense that for each $y \in \Gamma$, $\bar{\varphi}(y) \cap \Gamma \neq \emptyset$. Therefore we can define recursively a sequence $\{y_n\} \subseteq \Gamma$, setting $y_0 = z$, and picking for each $n \in \mathbb{N}$, $y_{n+1} \in \bar{\varphi}(y_n) \cap \Gamma$. Let y be a sublimit of $\{y_n\}$. Then $y \in \Gamma \cap R(\varphi) \subseteq \bar{U} \cap R(\varphi)$, which proves that $x \in W_R^u(\bar{U} \cap R(\varphi))$. \square

Let $\varphi, \psi \in \mathcal{O}(X)$. We say that φ is *combinatorially equivalent* to ψ , and write $\varphi \bowtie \psi$, if and only if the permutations π_φ and π_ψ are conjugated, that is, there is a bijective map $h : \Sigma_{\text{final}}^\Omega(\varphi) \rightarrow \Sigma_{\text{final}}^\Omega(\psi)$ such that the following diagram commutes:

$$\begin{array}{ccc} \Sigma_{\text{final}}^\Omega(\varphi) & \xrightarrow{h} & \Sigma_{\text{final}}^\Omega(\psi) \\ \pi_\varphi \downarrow & & \downarrow \pi_\psi \\ \Sigma_{\text{final}}^\Omega(\varphi) & \xrightarrow{h} & \Sigma_{\text{final}}^\Omega(\psi) \end{array} .$$

Given a topological subsemigroup $\mathcal{O}_1 \subseteq \mathcal{O}(X)$, we say that $\varphi \in \mathcal{O}(X)$ is *combinatorially stable in \mathcal{O}_1* if and only if there is a neighbourhood \mathcal{U} of φ in \mathcal{O}_1 such that all $\psi \in \mathcal{U}$ are combinatorially equivalent to φ .

Given $\varphi, \psi \in \mathcal{O}(X)$ we define the following multi-valued mappings:

If $\varphi \leq \psi$ let

$$\tau_{\varphi, \psi} : \Lambda_{\text{final}}^\Omega(\psi) \rightarrow \mathcal{P}(\Lambda_{\text{final}}^\Omega(\varphi)) \quad \text{and} \quad \theta_{\varphi, \psi} : \Sigma_{\text{final}}^\Omega(\psi) \rightarrow \mathcal{P}(\Sigma_{\text{final}}^\Omega(\varphi)) ,$$

be the maps that to each Ω -final class, resp. Ω -final component, C of ψ associate the set of Ω -final classes, resp. Ω -final components, C' of φ contained in C .

If $\varphi \prec \psi$ let

$$\tau_{\varphi, \psi}^* : \Lambda_{\text{final}}^\Omega(\psi) \rightarrow \mathcal{P}(\Lambda_{\text{final}}^R(\varphi)) \quad \text{and} \quad \theta_{\varphi, \psi}^* : \Sigma_{\text{final}}^\Omega(\psi) \rightarrow \mathcal{P}(\Sigma_{\text{final}}^R(\varphi)) ,$$

be the maps that to each Ω -final class, resp. Ω -final component, C of ψ associate the set of R -final classes, resp. R -final components, A of φ contained in C .

In general, for any φ let

$$\tilde{\tau}_\varphi : \Lambda_{\text{final}}^R(\varphi) \rightarrow \mathcal{P}(\Lambda_{\text{final}}^\Omega(\varphi)) \quad \text{and} \quad \tilde{\theta}_\varphi : \Sigma_{\text{final}}^R(\varphi) \rightarrow \mathcal{P}(\Sigma_{\text{final}}^\Omega(\varphi)) ,$$

be the maps that to each R -final class, resp. R -final component, A of φ associate the set of Ω -final classes, resp. Ω -final components, C of φ contained in A .

Proposition 5.10. *Let $\varphi_1, \varphi_2, \varphi_3 \in \mathcal{O}(X)$.*

If $\varphi_1 \prec \varphi_2$ then $\theta_{\varphi_1, \varphi_2} = \tilde{\theta}_{\varphi_1} \circ \theta_{\varphi_1, \varphi_2}^*$ and $\tau_{\varphi_1, \varphi_2} = \tilde{\tau}_{\varphi_1} \circ \tau_{\varphi_1, \varphi_2}^*$.
If $\varphi_1 \leq \varphi_2 \leq \varphi_3$ then $\theta_{\varphi_1, \varphi_3} = \theta_{\varphi_1, \varphi_2} \circ \theta_{\varphi_2, \varphi_3}$ and $\tau_{\varphi_1, \varphi_3} = \tau_{\varphi_1, \varphi_2} \circ \tau_{\varphi_2, \varphi_3}$.

Proposition 5.11. Let $\Theta : \Xi_1 \rightarrow \mathcal{P}(\Xi_2)$ stand for any of the multi-valued mappings, $\theta_{\varphi, \psi}$, $\theta_{\varphi, \psi}^*$, $\tilde{\theta}_{\varphi}$, $\tau_{\varphi, \psi}$, $\tau_{\varphi, \psi}^*$ or $\tilde{\tau}_{\varphi}$. Then for every $C \in \Xi_1$ and $A \in \Xi_2$, $\Theta(C) \neq \emptyset$, and $A \in \Theta(C) \Leftrightarrow A \subseteq C$. Furthermore, the following statements are equivalent

1. Θ is one-to-one from Ξ_1 onto Ξ_2 ,
2. Θ is single valued and surjective,
3. $|\Xi_1| = |\Xi_2|$.

When either one of the equivalent statements in the previous proposition holds we shall say that Θ is a *final equivalence*.

Proposition 5.12 (Combinatorics monotonicity). Let $\varphi, \psi \in \mathcal{O}(X)$.

1. $|\Lambda_{\text{final}}^{\Omega}(\varphi)| \geq |\Lambda_{\text{final}}^R(\varphi)|$ and $|\Sigma_{\text{final}}^{\Omega}(\varphi)| \geq |\Sigma_{\text{final}}^R(\varphi)|$,
2. If $\varphi \leq \psi$, then $|\Lambda_{\text{final}}^{\Omega}(\varphi)| \geq |\Lambda_{\text{final}}^{\Omega}(\psi)|$ and $|\Sigma_{\text{final}}^{\Omega}(\varphi)| \geq |\Sigma_{\text{final}}^{\Omega}(\psi)|$,
3. If $\varphi \prec \psi$, then $|\Lambda_{\text{final}}^R(\varphi)| \geq |\Lambda_{\text{final}}^R(\psi)|$ and $|\Sigma_{\text{final}}^R(\varphi)| \geq |\Sigma_{\text{final}}^R(\psi)|$,
4. Let Θ be any of the mappings, $\theta_{\varphi, \psi}$, $\theta_{\varphi, \psi}^*$ or $\tilde{\theta}_{\varphi}$. Given final components A and C , if $A \in \Theta(C)$ then the period of C divides the period of A .

Proposition 5.13. Given $\varphi, \psi \in \mathcal{O}(X)$ such that $\varphi \leq \psi$, the following statements are equivalent:

1. φ is combinatorially equivalent to ψ .
2. $|\Sigma_{\text{final}}^{\Omega}(\varphi)| = |\Sigma_{\text{final}}^{\Omega}(\psi)|$.
3. $|\Lambda_{\text{final}}^{\Omega}(\varphi^d)| = |\Lambda_{\text{final}}^{\Omega}(\psi^d)|$, for all $d \in \mathbb{N}$.

Proof. Assume that $\varphi \leq \psi$ and take d to be a multiple of all periods of Ω -final classes of φ . Then all Ω -final classes of φ^d are connected and $\Sigma_{\text{final}}^\Omega(\varphi) = \Lambda_{\text{final}}^\Omega(\varphi^d)$. Since d is also a multiple of all periods of classes in $\Lambda_{\text{final}}^\Omega(\psi)$ we have $\Sigma_{\text{final}}^\Omega(\psi) = \Lambda_{\text{final}}^\Omega(\psi^d)$. These equalities show that 3. implies 2.. \square

Lemma 5.14 (Factorization lemma). *In each of the factorizations of proposition 5.10, the composition is a final equivalence if and only if both factors are final equivalences.*

Proposition 5.15. *Given $\varphi, \psi \in \mathcal{O}(X)$, such that $\varphi \leq \psi$, if $\theta_{\varphi, \psi}$ is a final equivalence then every map $\zeta \in \mathcal{O}(X)$ such that $\varphi \leq \zeta \leq \psi$ is combinatorially equivalent to both φ and ψ .*

Proposition 5.16 (Upper stability of final combinatorics). *Given $\varphi \in \mathcal{O}(X)$, there is some $\epsilon > 0$ such that $\theta_{\varphi, \varphi_\epsilon}^*$ is a final equivalence.*

Proposition 5.17 (Lower stability of final combinatorics). *Given $\varphi \in \mathcal{O}(X)$, there is some $\epsilon > 0$ such that $\theta_{\varphi_\epsilon, \varphi}$ is a final equivalence.*

If any of these statements holds for some $\epsilon_0 > 0$ then it also holds for all $0 < \epsilon < \epsilon_0$. This follows from the factorization lemma 5.14.

Lemma 5.18. *Every $\varphi \in \mathcal{O}(X)$ is combinatorially equivalent to $\widehat{\varphi}$.*

Proof. Given $\varphi \in \mathcal{O}(X)$, the open set $S(\varphi) = \bigcup_{n=1}^\infty \text{graph}(\varphi^n)$ is the graph of the pre-order \rightsquigarrow_φ . The intersection $E(\varphi) = S(\varphi) \cap S(\varphi)^{-1}$ is an open set which together with the diagonal $\Delta_X \subseteq X \times X$ gives the graph of the equivalence relation \leftrightarrow_φ . Finally, denoting by $\pi : X \times X \rightarrow X$ the projection onto the first factor, we have $\Omega(\varphi) = \pi(E(\varphi))$.

By proposition 4.3 it follows that $\text{graph} \varphi^n$ is dense in $\text{graph}(\widehat{\varphi})^n$, for each $n \geq 1$. Therefore $S(\varphi)$ is dense in $S(\widehat{\varphi})$, which implies that $E(\varphi)$ is dense in $E(\widehat{\varphi})$, and in turn this implies that $\Omega(\varphi)$ is dense in $\Omega(\widehat{\varphi})$. Given open sets $U_1, U_2 \subseteq \Omega(\varphi)$, since $S(\varphi)$ is dense in $S(\widehat{\varphi})$, $U_1 \rightsquigarrow_{\widehat{\varphi}} U_2$ implies $U_1 \rightsquigarrow_\varphi U_2$. This shows that the map $\tau_{\varphi, \widehat{\varphi}} : \Lambda_{\text{final}}^\Omega(\widehat{\varphi}) \rightarrow \mathcal{P}(\Lambda_{\text{final}}^\Omega(\varphi))$ is bijective.

Because of proposition 4.3, we have $\varphi^d \leq (\widehat{\varphi})^d \leq \widehat{(\varphi^d)}$ and applying the previous argument to φ^d we see that

$$|\Lambda_{\text{final}}^{\Omega}(\varphi^d)| = |\Lambda_{\text{final}}^{\Omega}((\widehat{\varphi})^d)| = |\Lambda_{\text{final}}^{\Omega}(\widehat{(\varphi^d)})|,$$

for every $d \in \mathbb{N}$. By proposition 5.13 this implies that φ and $\widehat{\varphi}$ are combinatorially equivalent. \square

Corollary 5.19. *All open maps in a class $[\varphi]$ of the quotient semigroup $\mathbb{O}(X)$ are combinatorially equivalent to each other.*

Let $\varphi \in \mathcal{O}(X)$. We shall call *thickness* of φ to the smallest volume (m -measure) of all its components in $\Sigma_{\text{final}}^{\Omega}(\varphi)$. We say that an open set $K \subseteq \Omega_{\text{final}}(\varphi)$ is a *final kernel* of φ if and only if there is a one-to-one correspondence $C \mapsto K_C$, between components $C \in \Sigma_{\text{final}}^{\Omega}(\varphi)$ and connected components K_C of K , such that $\overline{K_C} \subseteq C$ for every $C \in \Sigma_{\text{final}}^{\Omega}(\varphi)$. We say that K is a *final kernel with finite order N* if and only if K is a final kernel of φ , and furthermore

1. For each component $C \in \Sigma_{\text{final}}^{\Omega}(\varphi)$ of period d , the only connected component K_C of K contained in C satisfies $\overline{C \times K_C} \subseteq \text{graph } \varphi^{Nd}$.
2. For each $x \in X$, $\varphi^N(x)$ contains at least the closure of one of K 's connected components.

We shall call *thickness* of the final kernel K to the smallest volume of all of K 's connected components.

Lemma 5.20. *Given $\varphi \in \mathcal{O}(X)$, every final kernel K of φ is a final kernel with some finite order $N \in \mathbb{N}$. In particular, φ admits finite order final kernels, whose thickness is arbitrarily close to the thickness of φ .*

Proof. Let K be a final kernel of φ . Let $C \in \Sigma_{\text{final}}^{\Omega}(\varphi)$ be a connected component of period d and let K_C be the correspondent connected component of K . Fix $x_0 \in C$. There is $n \in \mathbb{N}$ such that $x_0 \in \varphi^n(x_0)$ and all such n 's are multiples of d . It is clear that if $x_0 \in \varphi^n(x_0)$ and $x_0 \in \varphi^m(x_0)$, then $x_0 \in \varphi^{n+m}(x_0)$. Consequently a simple number theory argument ensures that for some big enough n_0 and for all $n \geq n_0$, one has $x_0 \in$

$\varphi^{nd}(x_0)$. For any $(x, y) \in \overline{C} \times C$ there is $n \in \mathbb{N}$ such that $y \in \varphi^n(x)$ and all such n 's are multiples of d . Furthermore, because φ is open, the map $n(x, y) = \inf \{n : y \in \varphi^n(x)\}$ is upper semi-continuous in $(x, y) \in \overline{C} \times C$. Therefore, there exists $N_0 = N_0(K_C)$ such that if $(x, y) \in \overline{C} \times \overline{K_C}$ then $x_0 \in \varphi^k(x)$ and $y \in \varphi^l(x_0)$, for some $k, l < N_0$. Set N_C to be the integral part of $n_0 + 2N_0/d + 1$. Then for every $n \geq N_C$, one has $y \in \varphi^{nd}(x)$, that is, $\overline{C} \times \overline{K_C} \subseteq \text{graph } \varphi^{nd}$ for every $n \geq N_C$. Set $N_1 = \max\{N_C : C \in \Sigma_{\text{final}}^\Omega(\varphi)\}$. Then for every $N \geq N_1$ and for each component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$ of period d , one has $\overline{C} \times \overline{K_C} \subseteq \text{graph } \varphi^{Nd}$, which proves item 1. in the definition of final kernel of order N .

For any $x \in X$ define $n(x) = \inf \{n : \varphi^n(x) \cap \Omega_{\text{final}}(\varphi) \neq \emptyset\}$. The map $n : X \rightarrow \mathbb{N}$ is bounded because $X = W_\Omega^s(\Omega_{\text{final}}(\varphi))$, and is upper semi-continuous because φ is open. Furthermore, since $\Omega_{\text{final}}(\varphi)$ is φ -invariant, if for some n_0 one has $\varphi^{n_0}(x) \cap \Omega_{\text{final}}(\varphi) \neq \emptyset$, then $\varphi^n(x) \cap \Omega_{\text{final}}(\varphi) \neq \emptyset$ for every $n \geq n_0$. Altogether these facts ensure that there is $N_2 \in \mathbb{N}$ such that for each $x \in X$, $\varphi^{N_2}(x) \cap \Omega_{\text{final}}(\varphi) \neq \emptyset$. Therefore, for each $x \in X$ there is $C \in \Sigma_{\text{final}}^\Omega(\varphi)$ such that $\varphi^{N_2}(x) \cap C \neq \emptyset$. From above we know that $\overline{C} \times \overline{K_C} \subseteq \text{graph } \varphi^{N_1 d}$. Therefore, setting $N = N_1 d + N_2$ one has that for each $x \in X$, $\varphi^N(x)$ contains at least the closure of one of K 's connected components. This proves item 2. in the definition of final kernel of finite order N . Thus K is a final kernel of order N . \square

Proof. (of proposition 5.17) Take some final kernel $K \subseteq \Omega_{\text{final}}(\varphi)$ and choose $N \in \mathbb{N}$ such that K is a finite kernel of order N . It is clear that for small enough $\epsilon > 0$, $K \subseteq \Omega_{\text{final}}(\varphi_\epsilon^\circ)$ is also a final kernel of the same order N for the map φ_ϵ° . We claim that for such ϵ , $\theta_{\varphi_\epsilon^\circ, \varphi}$ is a final equivalence. Indeed, item 1. in the definition of final kernel implies that for a given component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$ there is at most one component $C_\epsilon \in \Sigma_{\text{final}}^\Omega(\varphi_\epsilon^\circ)$ contained in C : the component that contains K_C . On the other hand, item 2. of the same definition implies that any final component $C_\epsilon \in \Sigma_{\text{final}}^\Omega(\varphi_\epsilon^\circ)$ must contain one of K 's components and, therefore, be contained inside a final component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$. These two facts together show that $\theta_{\varphi_\epsilon^\circ, \varphi}$ is a final equivalence. \square

Lemma 5.21. *Given $\varphi \in \mathcal{O}$, there is some $\epsilon > 0$ such that $\tau_{\varphi, \varphi_\epsilon^*}^*$ is a final equivalence.*

Proof. Since for any pair (A_1, A_2) of classes $A_1, A_2 \in \Lambda_{\text{final}}^R(\varphi)$, we have $A_1 \not\sim_\varphi A_2$, we can choose $\epsilon > 0$ small enough so that for all those pairs $A_1 \not\sim_{\varphi_\epsilon^*} A_2$.

We have to prove that any final class $A \in \Lambda_{\text{final}}^R(\varphi)$ is contained in some final class $C \in \Lambda_{\text{final}}^\Omega(\varphi_\epsilon^*)$. Given $A \in \Lambda_{\text{final}}^R(\varphi)$ clearly there is some $C \in \Lambda_{\text{final}}^\Omega(\varphi_\epsilon^*)$ such that

$A \rightsquigarrow_{\varphi_\epsilon^*} C$, and for this class C there is $A' \in \Lambda_{\text{final}}^R(\varphi)$ such that $A' \subseteq C$. Therefore $A \rightsquigarrow_{\varphi_\epsilon^*} A'$, which by ϵ' choice implies $A \rightarrow_\varphi A'$, and so, since A is a final class, $A = A'$.

Finally we have to see that given $C \in \Lambda_{\text{final}}^\Omega(\varphi_\epsilon^*)$ there is at most one class $A \in \Lambda_{\text{final}}^R(\varphi)$ contained in C . Assume $A_1, A_2 \in \Lambda_{\text{final}}^R(\varphi)$ are two final classes contained inside $C \in \Lambda_{\text{final}}^\Omega(\varphi_\epsilon^*)$. Then $A_1 \rightsquigarrow_{\varphi_\epsilon^*} A_2$ and $A_2 \rightsquigarrow_{\varphi_\epsilon^*} A_1$, which by ϵ' choice implies $A_1 \rightarrow_\varphi A_2$ and $A_2 \rightarrow_\varphi A_1$. This proves that $A_1 = A_2$. \square

Lemma 5.22. *There is $\epsilon_0 > 0$ such that $\{\Omega_{\text{final}}(\varphi_\epsilon^*)\}_{0 < \epsilon < \epsilon_0}$ is an outer approximation of $R_{\text{final}}(\varphi)$.*

Proof. Take $\epsilon_0 > 0$ given by lemma 5.21. Then for $0 < \epsilon < \epsilon_0$, $R_{\text{final}}(\varphi) \subseteq \Omega_{\text{final}}(\varphi_\epsilon^*)$. Since it is clear that $\{\Omega(\varphi_\epsilon^*)\}_{0 < \epsilon < \epsilon_0}$ is an outer approximation of $R(\varphi)$, the lemma follows. \square

Proof. (of proposition 5.16) Let $\gamma > 0$ be the minimum of all distances between components in $\Sigma_{\text{final}}^R(\varphi)$. Take $\epsilon > 0$ small enough so that $\tau_{\varphi, \varphi_\epsilon^*}^*$ is a final equivalence and $R_{\text{final}}(\varphi) \subseteq \Omega_{\text{final}}(\varphi_\epsilon^*) \subseteq B_{\gamma/2}(R_{\text{final}}(\varphi))$. It follows that each connected component of $R_{\text{final}}(\varphi)$ is contained inside a connected component of $\Omega_{\text{final}}(\varphi_\epsilon^*)$, and each connected component of $\Omega_{\text{final}}(\varphi_\epsilon^*)$ contains at most one connected component of $R_{\text{final}}(\varphi)$. Therefore $\theta_{\varphi, \varphi_\epsilon^*}^*$ is a final equivalence. \square

We say that $\varphi \in \mathcal{O}(X)$ satisfies the *combinatorial stability condition* if and only if $\tilde{\theta}_\varphi$ is a final equivalence, or equivalently iff $|\Lambda_{\text{final}}^\Omega(\varphi)| = |\Lambda_{\text{final}}^R(\varphi)|$ and $|\Sigma_{\text{final}}^\Omega(\varphi)| = |\Sigma_{\text{final}}^R(\varphi)|$.

Theorem 5.3 (Stability characterization). *Let \mathcal{O}_1 be any topological semigroup of open maps. For any $\varphi \in \mathcal{O}_1$, φ is combinatorially stable in \mathcal{O}_1 if and only if φ satisfies the combinatorial stability condition.*

Proof. In view of corollary 5.19 it's enough proving this theorem for the quotient semigroup $\mathbb{O}_1 = \mathcal{O}_1 / \sim$.

Given $\varphi \in \mathbb{O}_1$ assume that $\tilde{\theta}_\varphi$ is a final equivalence. Let $K \subseteq \Omega_{\text{final}}(\varphi)$ be some final kernel with finite order $N \in \mathbb{N}$ and positive thickness $c > 0$. Let K_1, \dots, K_p be the connected components of K , and C_1, \dots, C_p be the correspondent components in $\Sigma_{\text{final}}^\Omega(\varphi)$. Take $0 < \epsilon < c/3$ as in proposition 5.16. Find an open cover of X ,

$\{U_i : 1 \leq i \leq p\}$, such that $\overline{U_i \times K_i} \subseteq \overline{\text{graph}(\varphi^N)}$ for every $i = 1, \dots, p$. Notice that, since K is a final kernel of order N , we have $\overline{C_i \times K_i} \subseteq \overline{\text{graph}(\varphi^{Nd_i})}$, where d_i is the period of C_i . Then choose some neighbourhood \mathcal{N} of φ in \mathbb{O}_1 such that for every $\psi \in \mathcal{N}$, $\rho_H(\psi, \varphi) < \epsilon$, $m(K_i \setminus \psi^N(x)) < \epsilon$ if $x \in \overline{U_i}$, and $m(K_i \setminus \psi^{Nd_i}(x)) < \epsilon$ when $x \in \overline{C_i}$, for $i = 1, \dots, p$. Such a neighbourhood \mathcal{N} exists by definition 4.2. Take any ψ in \mathcal{N} . Inequality $\rho_H(\psi, \varphi) < \epsilon$ implies $\psi \leq \varphi_\epsilon^*$, and so we may consider the mapping $\theta_\epsilon = \theta_{\psi, \varphi_\epsilon^*} : \Sigma_{\text{final}}^\Omega(\varphi_\epsilon^*) \rightarrow \mathcal{P}(\Sigma_{\text{final}}^\Omega(\psi))$. Let us show now that θ_ϵ is a final equivalence, which, in view of ϵ choice, proves that ψ is combinatorially equivalent to φ . Given $B \in \Sigma_{\text{final}}^\Omega(\psi)$, pick an index $i = 1, \dots, p$ such that $B \cap U_i \neq \emptyset$ and take $x \in B \cap U_i$. Because $m(K_i \setminus \psi^N(x)) < \epsilon < m(K_i)/3$, we have $m(K_i \cap \psi^N(x)) > 2m(K_i)/3$, which implies $\psi^N(x) \cap K_i \neq \emptyset$. So there is some $z \in K_i \subseteq \Omega_{\text{final}}(\varphi_\epsilon^*)$ such that $B \rightsquigarrow_\psi z$. Thus, since $\psi \leq \varphi_\epsilon^*$, $B \subseteq \Omega_{\text{final}}(\varphi_\epsilon^*)$. This proves, denoting by C the Ω -final component of φ_ϵ^* containing z , that $B \in \theta_\epsilon((\pi_{\varphi_\epsilon^*})^{-N}(C))$. Therefore θ_ϵ is surjective. Let now $B_1, B_2 \in \Sigma_{\text{final}}^\Omega(\psi)$ be two components such that $B_1, B_2 \in \theta_\epsilon(\tilde{C}_j)$ where for some $j = 1, \dots, p$, $\tilde{C}_j \in \Sigma_{\text{final}}^\Omega(\varphi_\epsilon^*)$ is the unique component that contains C_j . Since $K_j \subseteq C_j \subseteq \tilde{C}_j$, arguing as before, for each $i = 1, 2$, there is some point $x_i \in K_j$ such that $B_i \rightsquigarrow_\psi x_i$. Since these are final components, we may choose the points x_i so that $x_i \in B_i$. Because $x_1, x_2 \in K_j \subseteq C_j$, by the choice of neighbourhood \mathcal{N} , we have

$$m(K_j \setminus \psi^{Nd_j}(x_1)) < m(K_j)/3 \quad \text{and} \quad m(K_j \setminus \psi^{Nd_j}(x_2)) < m(K_j)/3,$$

and so

$$m(K_j \cap \psi^{Nd_j}(x_1) \cap \psi^{Nd_j}(x_2)) \geq m(K_j)/3,$$

which implies that $\psi^{Nd}(x_1) \cap \psi^{Nd}(x_2) \neq \emptyset$. Since B_1 and B_2 are final components, it follows that $B_1 = B_2$. This proves that θ_ϵ is single valued and, therefore, a final equivalence.

Assume now that φ is \mathbb{O}_1 -combinatorially stable and let \mathcal{N} be a neighbourhood of φ in \mathbb{O}_1 where all open maps are combinatorially equivalent. Take $\epsilon > 0$ as in proposition 5.16. By item (b) of definition 3.3, used in definition 4.2, there is an outer approximation $\{\varphi_n\}$ of φ such that $\lim_{n \rightarrow \infty} \varphi_n = \varphi$. Then for all large enough n , $\varphi_n \in \mathcal{N}$ and $\varphi_n \prec \varphi_\epsilon^*$. Therefore $\theta_{\varphi, \varphi_n}$ is a final equivalence, because φ and φ_n are combinatorially equivalent; and $\theta_{\varphi, \varphi_n}^*$ is also a final equivalence, by the choice of ϵ in proposition 5.16. Since $\theta_{\varphi, \varphi_n} = \theta_\varphi \circ \theta_{\varphi, \varphi_n}^*$, the Factorization lemma 5.10 implies that $\tilde{\theta}_\varphi$ is a final equivalence. \square

Theorem 5.4 (Genericity of combinatorial stability). *Let \mathcal{O}_1 be any topological semigroup of open maps. The set of \mathcal{O}_1 -combinatorially stable maps is open and dense in the semigroup \mathcal{O}_1 .*

Proof. By definition, combinatorial stability is an open property. Thus it is enough to prove density. Consider any open map $\varphi \in \mathcal{O}_1$. Take $\epsilon > 0$ as in proposition 5.16. By item (b) of definition 3.3, used in definition 4.2, there is an outer approximation $\{\varphi_n\}$ of φ such that $\lim_{n \rightarrow \infty} \varphi_n = \varphi$. Then for all large enough n , $\varphi_n \prec \varphi_\epsilon^*$, and since $\theta_{\varphi, \varphi_\epsilon^*}^* = \theta_{\varphi, \varphi_n}^* \circ \tilde{\theta}_{\varphi_n} \circ \theta_{\varphi_n, \varphi_\epsilon^*}^*$, the Factorization lemma 5.10 implies that $\tilde{\theta}_{\varphi_n}$ is a final equivalence. Thus, φ is the limit of the combinatorially stable open maps φ_n . \square

We say that an Ω -final class of φ is *unstable* when it is contained inside an R -class of φ which is not final.

We say that two or more Ω -final classes C_1, \dots, C_k of φ *collapse together* when they are contained in the same R -final class of φ .

We say that *the period of an Ω -final class C of φ is collapsing* when it is contained in one R -final class whose period is less than C 's period.

The following proposition describes the *bifurcation set* of φ in a topological semigroup of open maps. It's just an obvious reformulation of theorem 5.3.

Proposition 5.23. *Let \mathcal{O}_1 be any topological semigroup of open maps. Given $\varphi \in \mathcal{O}_1$, φ is not \mathcal{O}_1 -combinatorially stable if and only if one of the following alternatives hold:*

- (a) *One Ω -final class of φ is unstable,*
- (b) *Two or more Ω -final classes of φ collapse together,*
- (c) *The period of one Ω -final class C of φ is collapsing.*

Recall that a space is called a *Baire space* if and only if any countable intersection of open and dense subsets is still dense. By Baire Theorem, any complete metric space is a Baire space. In particular, compact metric spaces are Baire spaces. It is also straightforward to check that an inductive limit of Baire spaces is again a Baire space. Therefore since the topological semigroup $\mathbb{O}_{\text{Lip}}^r$ is the inductive limit of compact metric

spaces, it is a Baire space. On Baire spaces, a countable intersection of open and dense subsets is called a *residual* subset.

Let us say that an open map $\varphi \in \mathcal{O}(X)$ satisfies the *topological stability condition* if and only if φ satisfies the combinatorial stability condition, $\overline{\Omega_{\text{final}}(\varphi)} = R_{\text{final}}(\varphi)$ and both these sets are Jordan measurable. Set then

$$\mathcal{O}_{\text{stable}} = \{ \varphi \in \mathcal{O}(X) : \varphi \text{ satisfies the topological stability condition} \}.$$

We say that a closed set $K \subseteq X$ has *regular boundary* if $\partial K = \partial(K^\circ) \Leftrightarrow K = \overline{K^\circ}$.

Lemma 5.24. *For every $\varphi \in \mathcal{O}_{\text{Cont}}(X)$, the closed set $R_{\text{final}}(\varphi)$ has regular boundary.*

Proof. It is obvious that the closure of an open set is always a closed set with regular boundary. Then for any open map $\varphi \in \mathcal{O}_{\text{Cont}}(X)$, and any $x \in X$, $\overline{\varphi(x)} = \overline{\varphi(x)}$ has regular boundary. The equality holds because φ is continuous. Thus, for each class $C \in \Lambda_{\text{final}}^R(\varphi)$, $C = \overline{\varphi(C)} = \overline{\varphi(C)}$, which proves that C , being the closure of the open set $\varphi(C)$, has regular boundary. Since $\Lambda_{\text{final}}^R(\varphi)$ is finite, $R_{\text{final}}(\varphi)$ is a finite disjoint union of closed sets with regular boundary, and therefore has itself regular boundary. \square

Lemma 5.25. *Given $\varphi \in \mathcal{O}_{\text{Cont}}(X)$, let $\{\varphi_\epsilon^*\}_{0 < \epsilon < \epsilon_0}$ be any outer approximation of φ by continuous open maps. Then the set $\{\epsilon \in]0, \epsilon_0[: \varphi_\epsilon^* \notin \mathcal{O}_{\text{stable}}\}$ is at most countable. In particular, in any topological semigroup of open maps $\mathcal{O}_1 \subseteq \mathcal{O}_{\text{Cont}}(X)$, the set $\mathcal{O}_1 \cap \mathcal{O}_{\text{stable}}$ is dense in \mathcal{O}_1 .*

A similar result holds for inner approximations.

Proof. Let $A_\epsilon = R_{\text{final}}(\varphi_\epsilon^*)^\circ \setminus \overline{\Omega_{\text{final}}(\varphi_\epsilon^*)}$ and $B_\epsilon = R_{\text{final}}(\varphi_\epsilon^*) \setminus \Omega_{\text{final}}(\varphi_\epsilon^*)$ for each $0 < \epsilon < \epsilon_0$. Then $\{A_\epsilon\}_{\epsilon > 0}$ is a family of pairwise disjoint open sets, and $\{B_\epsilon\}_{\epsilon > 0}$ is a family of pairwise disjoint closed sets. By monotonicity of the combinatorics, see proposition 5.12, the set $\Xi_0 = \{\epsilon \in]0, \epsilon_0[: \tilde{\theta}_{\varphi_\epsilon^*} \text{ is not a final equivalence}\}$ is at most finite. Since the topology of X admits countable open sub-basis, the set $\Xi_1 = \{\epsilon \in]0, \epsilon_0[: A_\epsilon \neq \emptyset\}$ is at most countable. Because $\sum_{0 < \epsilon < \epsilon_0} m(B_\epsilon) \leq m(X) = 1$, the set $\Xi_2 = \{\epsilon \in]0, \epsilon_0[: m(B_\epsilon) > 0\}$ is, once more, at most countable. Finally, it should be clear that the set $\{\epsilon \in]0, \epsilon_0[: \varphi_\epsilon^* \notin \mathcal{O}_{\text{stable}}\}$ is covered by the countable union $\Xi_0 \cup \Xi_1 \cup \Xi_2$. At this point we use the previous lemma to ensure that $A_\epsilon = \emptyset$ implies $R_{\text{final}}(\varphi_\epsilon^*) = \overline{\Omega_{\text{final}}(\varphi_\epsilon^*)}$. \square

Theorem 5.5 (Continuity of the Attractors). *Let \mathbb{O}_1 be any topological semigroup of open maps.*

1. *The mappings $\varphi \mapsto \Omega_{\text{final}}(\varphi)$ from \mathbb{O}_1 into $(\mathbb{U}(X), \rho_H)$, and $\varphi \mapsto m(\Omega_{\text{final}}(\varphi))$ from \mathbb{O}_1 into \mathbb{R} are lower semi-continuous.*
2. *The mappings $\varphi \mapsto R_{\text{final}}(\varphi)$ from \mathbb{O}_1 into $(\mathbb{K}(X), \rho_H)$, and $\varphi \mapsto m(R_{\text{final}}(\varphi))$ from \mathbb{O}_1 into \mathbb{R} are upper semi-continuous.*
3. *The four mappings on \mathbb{O}_1 , $\varphi \mapsto \Omega_{\text{final}}(\varphi)$, $\varphi \mapsto R_{\text{final}}(\varphi)$, $\varphi \mapsto m(R_{\text{final}}(\varphi))$ and $\varphi \mapsto m(\Omega_{\text{final}}(\varphi))$ are continuous at all $\varphi \in \mathcal{O}_1 \cap \mathcal{O}_{\text{stable}}(X)$.*
4. *The final components in $\Sigma_{\text{final}}^\Omega(\varphi)$ depend continuously on φ at those open maps in $\mathbb{O}_1 \cap \mathbb{O}_{\text{stable}}$, i.e. given $\varphi \in \mathbb{O}_1 \cap \mathbb{O}_{\text{stable}}$, there is a neighbourhood \mathcal{N} of φ in \mathbb{O}_1 , and for each $\psi \in \mathcal{N}$ there is a conjugation $h_{\varphi, \psi} : \Sigma_{\text{final}}^\Omega(\varphi) \rightarrow \Sigma_{\text{final}}^\Omega(\psi)$ between π_φ and π_ψ such that $\lim \rho_H(h_{\varphi, \psi}(C), C) = 0$, as ψ tends to φ , for every component $C \in \Sigma_{\text{final}}^\Omega(\varphi)$.*

Proof. Given $\varphi \in \mathbb{O}_1$ let K be a final kernel of φ such that $B_\epsilon(K) \supseteq \Omega_{\text{final}}(\varphi)$ and $m(K) \geq m(\Omega_{\text{final}}(\varphi)) - \epsilon$. Let K_1, K_2, \dots, K_p be the connected components of K , and C_1, C_2, \dots, C_p be the corresponding components in $\Sigma_{\text{final}}^\Omega(\varphi)$. Denote by d_i the period of component C_i , for $i = 1, \dots, p$. Choose $N \in \mathbb{N}$ such that kernel K has order N . Arguing as in the proof of theorem 5.3, we can find a neighbourhood \mathcal{N} of φ such that for each $\psi \in \mathcal{N}$ and $i = 1, \dots, p$, $K_i \cap \Omega_{\text{final}}(\psi) \neq \emptyset$. By proposition 4.8 we can make \mathcal{N} smaller so that for every $\psi \in \mathcal{N}$, every $i = 1, \dots, p$, and each $x \in K_i$, one has $B_\epsilon(\psi^{N d_i}(x)) \supseteq \varphi^{N d_i}(x)$. By definition 4.2 we make \mathcal{N} even smaller so that for every $\psi \in \mathcal{N}$, every $i = 1, \dots, p$, and each $x \in C_i$, one has $m(K_i \setminus \psi^{N d_i}(x)) < \epsilon/p$. Now pick any $\psi \in \mathcal{N}$. For each $i = 1, \dots, p$ choose $x_i \in K_i \cap \Omega_{\text{final}}(\psi)$. Then

$$B_\epsilon(\Omega_{\text{final}}(\psi)) \supseteq \cup_{i=1}^p B_\epsilon(\psi^{N d_i}(x_i)) \supseteq \cup_{i=1}^p \varphi^{N d_i}(x_i) \supseteq \cup_{i=1}^p K_i = K .$$

And so

$$B_{2\epsilon}(\Omega_{\text{final}}(\psi)) \supseteq B_\epsilon(K) \supseteq \Omega_{\text{final}}(\varphi) .$$

Analogously

$$\begin{aligned}
m(\Omega_{\text{final}}(\psi)) &\geq \sum_{i=1}^p m(\psi^{N d_i}(x_i) \cap K_i) \\
&= \sum_{i=1}^p m(K_i) - m(K_i \setminus \psi^{N d_i}(x_i)) \\
&> m(K) - \sum_{i=1}^p \epsilon/p = m(K) - \epsilon \\
&\geq m(\Omega_{\text{final}}(\varphi)) - 2\epsilon .
\end{aligned}$$

This proves the lower continuity of both mappings and establishes item 1. The upper continuity in item 2. follows from lemma 5.22. Item 3. follows easily from items 1. and 2. Finally, item 4. follows from 3. \square

Theorem 5.6. *Given any topological semigroup of open maps $\mathbb{O}_1 \subseteq \mathbb{O}_{\text{Cont}}$, the set $\mathbb{O}_1 \cap \mathbb{O}_{\text{stable}}$ is a countable intersection of open and dense subsets of \mathbb{O}_1 . In particular, for each $r \geq 0$, $\mathbb{O}_{\text{stable}} \cap \mathbb{O}_{\text{Lip}}^r$ is residual in $\mathbb{O}_{\text{Lip}}^r$.*

Proof. For each $\epsilon > 0$ set

$$\mathbb{O}_1^\epsilon = \{ \varphi \in \mathbb{O}_1 : B_\epsilon(\Omega_{\text{final}}(\varphi)) \supseteq R_{\text{final}}(\varphi) \text{ and } m(\Omega_{\text{final}}(\varphi)) \geq m(R_{\text{final}}(\varphi)) - \epsilon \}$$

By lemma 5.25, \mathbb{O}_1^ϵ is dense in \mathbb{O}_1 . Theorem 5.5 proves that each \mathbb{O}_1^ϵ is an open set in \mathbb{O}_1 . Therefore $\mathbb{O}_1 \cap \mathbb{O}_{\text{stable}} = \bigcap_{n=1}^\infty \mathbb{O}_1^{\frac{1}{n}}$ is a countable intersection of open and dense subsets of \mathbb{O}_1 . \square

6 Combinatorial Stability of Continuous Maps

Throughout this section we shall denote by $\mathcal{C}^0(X)$ the space of all continuous mappings $f : X \rightarrow X$, with the usual C^0 topology of uniform convergence. This topology is defined by the metric

$$d_H(f, g) = \max_{x \in X} d(f(x), g(x)) = \sup_{x \in X} \rho_H(\{f(x)\}, \{g(x)\}) ,$$

corresponding to distance between point-set maps introduced in 4.3. Let $f \in \mathcal{C}^0(X)$. Given $\epsilon > 0$, f_ϵ^* will denote the ϵ -explosion of f , which is defined by $\text{graph}(f_\epsilon^*) = B_\epsilon(\text{graph } f)$. Any sequence $x_0, x_1, \dots, x_n \in X$ such that $x_i \in f_\epsilon^*(x_{i-1})$ for $i = 1, \dots, n$ will be called an ϵ -pseudo orbit of f , and we shall say that x_n is an ϵ -pseudo iterate of the state x_0 . If y is an ϵ -pseudo iterate of x for every $\epsilon > 0$, y is said a pseudo iterate of x , and we shall write $x \rightarrow_f y$. Clearly \rightarrow_f is a transitive closed relation. Given subsets $A, B \subseteq X$, we shall write $A \rightarrow_f B$ when $x \rightarrow_f y$ for some $x \in A$ and some $y \in B$.

As usual, a subset $A \subseteq X$ is said to be f -invariant iff $f(A) \subseteq A$. An invariant set $A \subseteq X$ is called *chain transitive* iff $x \rightarrow_f y$ for every $x, y \in A$. We shall say that $A \subseteq X$ is an *acyclic attractor* iff A is a compact connected, f -invariant, chain transitive set and admits an *isolating* open neighbourhood $U \supseteq A$, in the sense that

$$\overline{f(U)} \subseteq U \quad \text{and} \quad A = \bigcap_{n=1}^{\infty} f^n(U).$$

When the set A splits as a disjoint union of d compact connected sets,

$$A = A_0 \cup f(A_0) \cup \dots \cup f^d(A_0),$$

such that A_0 is an acyclic attractor for f^d we say that A is a *periodic attractor* of period d . We define the *basin of attraction* of an attractor A , resp. the *pseudo basin of attraction*, as $W^s(A, f) = \bigcup_{n=0}^{\infty} f^{-n}(A)$, resp. $W_R^s(A, f) = \{x \in X : x \rightarrow_f A\}$. Finally, we say that the map f has *finitely many attractors* if and only if there is a finite number $p \in \mathbb{N}$ of periodic attractors A_1, \dots, A_p whose pseudo-basins of attraction cover the whole manifold, $X = \bigcup_{i=1}^p W_R^s(A_i, f)$.

We define the *chain recurrent set* of f , and denote it by $R(f)$, as the set of all states $x \in X$ such that x is a pseudo iterate of x , that is,

$$R(f) = \{x \in X : x \rightarrow_f x\}.$$

This set is closed in X , and the closed relation \rightarrow_f is a preorder on $R(f)$. Elements in $R(f)$ will be called *chain recurrent states*. Off course all points of an attractor are chain recurrent. The recurrent set $R(f)$ can be decomposed into classes defined by the following equivalence relation: $x \rightleftharpoons_f y \Leftrightarrow x \rightarrow_f y$ and $y \rightarrow_f x$, for any points $x, y \in R(f)$. Since \rightarrow_f is a preorder on $R(f)$, \rightleftharpoons_f is an equivalence relation. The

equivalence classes of recurrent states will be called *R-classes*. Clearly *R-classes* are closed subsets of X . As in the previous section, $\Lambda^R(f)$ shall denote the set of all *R-classes* of f . The relation \rightarrow_f is a partial ordering on $\Lambda^R(f)$. The maximal elements in $\Lambda^R(f)$ under \rightarrow_f will be called *R-final classes*, i.e. $D \in \Lambda^R(f)$ is an *R-final class* if and only if for all $C \in \Lambda^R(f)$, $D \rightarrow_f C$ implies $C = D$. As before, $\Lambda_{\text{final}}^R(f)$ will denote the set of all *R-final classes* of f . A chain recurrent state x will be called a *final chain recurrent state* if every pseudo iterate of x still has some pseudo iterate which comes back to x , that is,

$$R_{\text{final}}(f) = \{ x \in R(f) : \text{for every } y \in X, x \rightarrow_f y \Rightarrow y \rightarrow_f x \},$$

where we denote by $R_{\text{final}}(f)$ the set of all final chain recurrent states. We shall call *final chain recurrent set* to $R_{\text{final}}(f)$. This set is, in some sense, the union of system attractors. Next proposition clears the relation between attractors on one hand, and *R-final classes* and $R_{\text{final}}(f)$ on the other hand.

Proposition 6.1. *The set $R_{\text{final}}(f)$ has the following characterization:*

1. $R_{\text{final}}(f)$ is the union of all *R-final classes* $C \in \Lambda_{\text{final}}^R(f)$, and each such class is fully f -invariant, i.e. $f(C) = C$.
2. The attractors of f are the isolated *R-final classes* with finitely many connected components. More precisely, a subset $A \subseteq X$ is a periodic attractor of period d if and only if $A \in \Lambda_{\text{final}}^R(f)$, A has d connected components and there is an open set $U \subseteq X$ such that $U \cap R_{\text{final}}(f) = A$.
3. Each attractor admits a fundamental system of isolating neighbourhoods.

Proof. Item 1. follows by the same argument used in 5.6.

It is clear that an attractor of period d is an isolated *R-final class* with d connected components. Conversely, take $A \in \Lambda_{\text{final}}^R(f)$ with d connected components, isolated in $R_{\text{final}}(f)$. Then, one easily checks, the connected components A_0, A_1, \dots, A_{d-1} of A are cyclically permuted by f . Let $D = R_{\text{final}}(f) \setminus A$. Then D is closed since A is isolated. Choose $\gamma > 0$ such that the open balls $B_\gamma(D)$, and $B_\gamma(A_i)$, for $i = 0, 1, \dots, d-1$, are pairwise disjoint. Because $\{\Omega(f_\epsilon^*)\}_{\epsilon > 0}$ is an outer approximation of $R(f)$ we can take $\epsilon_0 > 0$ such that $\Omega(f_\epsilon^*) \subseteq B_\gamma(R(f))$, for all $0 < \epsilon < \epsilon_0$. Since $A \not\rightarrow_f D$ we can make $\epsilon_0 > 0$ smaller, but positive, so that $A \not\rightarrow_{f_\epsilon^*} D$ for every $0 < \epsilon < \epsilon_0$. Let U_i^ϵ be the connected component of $\Omega(f_\epsilon^*)$ that contains A_i ($0 \leq i \leq d-1$). Of course for

$0 < \epsilon < \epsilon_0$, U_i^ϵ is an Ω -final class in $\Sigma_{\text{final}}^\Omega(f_\epsilon^*)$. Then $A_i \subseteq U_i^\epsilon \subseteq B_\gamma(A_i)$, which implies that $A_i \subseteq \bigcap_{\epsilon>0} U_i^\epsilon \subseteq B_\gamma(A_i) \cap R(f)$. On the other hand the fact that each U_i^ϵ is a final class easily implies that the nested intersection $\bigcap_{\epsilon>0} U_i^\epsilon$ is contained in $B_\gamma(A_i) \cap R_{\text{final}}(f) = A_i$. Therefore $\{U_i^\epsilon\}_{\epsilon>0}$ is an outer approximation of A_i . Because U_i^ϵ is $(f_\epsilon^*)^d$ -fully invariant it follows that

$$\overline{f^d(U_i^\epsilon)} \subseteq (f_\epsilon^*)^d(U_i^\epsilon) = U_i^\epsilon .$$

Therefore, $\{U_i^\epsilon\}_{\epsilon>0}$ is a fundamental system of isolating neighbourhoods for A_i . Clearly

$$A_i = f^d(A_i) \subseteq \bigcap_{n=1}^{\infty} f^{nd}(U_i^\epsilon) .$$

On the other hand, for any $0 < \delta < \epsilon_1 < \epsilon$,

$$A_i \subseteq \bigcap_{n=1}^{\infty} f^{nd}(U_i^\epsilon) \subseteq \bigcap_{n=1}^{\infty} (f_\delta^*)^{nd}(U_i^\epsilon) \subseteq U_i^{\epsilon_1} ,$$

where last inclusion follows by proposition 5.8. Since $0 < \epsilon_1 < \epsilon$ is arbitrary small, it follows that $A_i = \bigcap_{n=1}^{\infty} f^{nd}(U_i^\epsilon)$ is a cyclic attractor of period d . \square

Lemma 6.2. *Given $f \in \mathcal{C}^0(X)$,*

$$R_{\text{final}}(f) = \overline{\lim_{\epsilon \rightarrow 0^+} \Omega_{\text{final}}(f_\epsilon^*)} := \overline{\bigcap_{\delta>0} \bigcup_{0<\epsilon<\delta} \Omega_{\text{final}}(f_\epsilon^*)} .$$

Proof. It is clear that $\{\Omega(f_\epsilon^*)\}_{\epsilon>0}$ is an outer approximation of $R(f)$. Therefore $\overline{\lim_{\epsilon \rightarrow 0^+} \Omega_{\text{final}}(f_\epsilon^*)} \subseteq R(f)$. Given $x \in \overline{\lim_{\epsilon \rightarrow 0^+} \Omega_{\text{final}}(f_\epsilon^*)}$, let $\{\epsilon_n\}$ be a sequence of positive numbers, decreasing to 0, such that for every $n \in \mathbb{N}$, $x \in \Omega_{\text{final}}(f_{\epsilon_n}^*)$. Let us prove that x is a final chain recurrent state. Assume $x \rightarrow_f y$. Then for all large enough $n \in \mathbb{N}$, $x \rightsquigarrow_{f_{\epsilon_n}^*} y$, which implies, because x is a final recurrent state for $f_{\epsilon_n}^*$, that $y \rightsquigarrow_{f_{\epsilon_n}^*} x$. Since $\epsilon_n > 0$ is arbitrary small, it follows that $y \rightarrow_f x$. Thus $x \in R_{\text{final}}(f)$.

Now, assume that $x \notin \overline{\lim_{\epsilon \rightarrow 0^+} \Omega_{\text{final}}(f_\epsilon^*)}$, and let us prove that $x \notin R_{\text{final}}(f)$. For some $\delta > 0$, $x \notin A_\delta = \bigcup_{0<\epsilon<\delta} \Omega_{\text{final}}(f_\epsilon^*)$. By proposition 5.3 we can take, for each $0 < \epsilon < \delta$, $z_\epsilon \in \Omega_{\text{final}}(f_\epsilon^*) \cap W_\Omega^s(x, f_\epsilon^*)$. Let z be a sublimit of $\{z_\epsilon\}_{\epsilon>0}$. Then $x \rightarrow_f z$. On the other hand $z \not\rightsquigarrow_{f_\epsilon^*} x$, because $x \notin A_\delta$ for each $0 < \epsilon < \delta$. This proves that x is not a final state. Therefore $x \notin R_{\text{final}}(f)$. \square

Proposition 6.3. *Given $f \in \mathcal{C}^0(X)$, $X = W_R^u(R_{\text{final}}(f))$.*

Proof. Given $x \in X$, take for each $\epsilon > 0$ $z_\epsilon \in \Omega_{\text{final}}(f_\epsilon^*)$ such that $x \rightsquigarrow_{f_\epsilon^*} z_\epsilon$. Let z be a sublimit of $\{z_\epsilon\}_{\epsilon>0}$. Then by the previous proposition $z \in R_{\text{final}}(f)$. It is also clear that $x \rightarrow_f z$. Thus $x \in W_R^u(R_{\text{final}}(f))$. \square

A connected component of $R_{\text{final}}(f)$ will be called an *R-final component* of f . We denote by $\Sigma_{\text{final}}^R(f)$ the quotient set of all *R-final components* of f . Unlike the case of open maps, the set $\Sigma_{\text{final}}^R(f)$ may be infinite. Nevertheless, the mapping $f : X \rightarrow X$ induces an action on the quotient $\bar{\pi}_f : \Sigma_{\text{final}}^R(f) \rightarrow \Sigma_{\text{final}}^R(f)$, defined by $\bar{\pi}_f(C) = f(C)$,

We say that a continuous mapping $f : X \rightarrow X$ is *weakly combinatorially stable* if and only if there is $\epsilon > 0$ such that all $\varphi \in \mathcal{O}(X)$ satisfying

1. $f \prec \varphi$, i.e. $\text{graph}(f) \subseteq \text{graph}(\varphi)$, and
2. $\text{diam}(\varphi) = \sup_{x \in X} \text{diam}(\varphi(x)) < \epsilon$

are combinatorially equivalent to f . We say that φ is *combinatorially equivalent to f* iff there is a bijective map $h : \Sigma_{\text{final}}^\Omega(\varphi) \rightarrow \Sigma_{\text{final}}^R(f)$ which conjugates the action of π_φ on $\Sigma_{\text{final}}^\Omega(\varphi)$ with that of $\bar{\pi}_f$ on $\Sigma_{\text{final}}^R(f)$, i.e. $\bar{\pi}_f \circ h = h \circ \pi_\varphi$.

Theorem 6.1. *Given $f \in \mathcal{C}^0(X)$, the following statements are equivalent:*

1. f has finitely many attractors,
2. $\Sigma_{\text{final}}^R(f)$ is a finite set,
3. f is weakly combinatorially stable,

Proof. 2. \Rightarrow 1. Assume that $\Sigma_{\text{final}}^R(f)$ is finite. Then every class $C \in \Lambda_{\text{final}}^R(f)$ is isolated with finitely many connected components, and, by proposition 6.1, is a periodic attractor of f . By proposition 6.3, the pseudo-basins of these attractors cover all X . Therefore f has finitely many attractors.

1. \Rightarrow 2. Obvious.

2. \Rightarrow 3. Assume that $\Sigma_{\text{final}}^R(f)$ is finite. For each $\epsilon > 0$ we can define, as in the previous section, $\theta_{f, f_\epsilon^*}^* : \Sigma_{\text{final}}^\Omega(f_\epsilon^*) \rightarrow \mathcal{P}(\Sigma_{\text{final}}^R(f))$ to be the map that to each class $C \in \Sigma_{\text{final}}^\Omega(f_\epsilon^*)$ assigns the set of *R-final classes* in $\Sigma_{\text{final}}^R(f)$ contained in C . With our finiteness assumption we can prove, as in proposition 5.16, that for some $\epsilon > 0$, $\theta_{f, f_\epsilon^*}^*$ is a final equivalence. Using the factorization lemma 5.14 we see that $\theta_{\varphi, f_\epsilon^*}$ is a final

equivalence for each open map $\varphi \in \mathcal{O}(X)$ with $f \prec \varphi \prec f_\epsilon^*$. Therefore f is weakly combinatorially stable.

3. \Rightarrow 2. Clear. □

We mention two examples of continuous maps $f \in \mathcal{C}^0(X)$ which are not weakly combinatorially stable: First we have maps with infinitely many attractors. They appear associated to Newhouse phenomenon of persistent heteroclinic tangencies in some hyperbolic invariant set. These classes of maps are residual in open sets of systems, w.r.t. the C^2 topology. Second we have infinitely renormalizable maps. Such examples admit totally disconnected 'attractors' and, therefore, can't be weakly combinatorially stable. Off course weak combinatorial stability can not be C^2 generic, as Newhouse examples show. But according to Palis conjecture this weak combinatorial stability should be dense in every space $\mathcal{C}^r(X)$ with the C^r topology.

Theorem 6.2. *The set of maps $f \in \mathcal{C}^0(X)$ with finitely many attractors is dense in $\mathcal{C}^0(X)$.*

The rest of this section is devoted to prove this theorem. Given $f \in \mathcal{C}^0(X)$ and $\epsilon > 0$ we shall denote by f_ϵ^\bullet the open map defined by $f_\epsilon^\bullet(x) = B_\epsilon(f(x))$, where the left hand side stands for the geodesic ball of radius $\epsilon > 0$. Since we are dealing with continuous mappings, $\{f_\epsilon^\bullet\}_{\epsilon > 0}$ is an outer approximation of f . By convention we set $f_0^\bullet = f$. Comparing with the standard outer approximation note that $f \prec f_\epsilon^\bullet \leq f_\epsilon^*$.

Given a finite subset $Z \subseteq X$, we shall call *skeleton-chain* to any map $\sigma : Z \rightarrow \mathcal{P}(Z)$ such that for every $x \in Z$ there is some $n \geq 1$ with $\sigma^n(x) = \emptyset$. States $x \in Z$ such that $\sigma(x) = \emptyset$ will be called *terminal states*. Similarly, when $\sigma^{-1}(x) = \emptyset$, $x \in Z$ will be called a *source state*. Clearly, a skeleton-chain is essentially a directed graph with no cycles. Consider open maps $\varphi, \psi \in \mathcal{O}(X)$. When a skeleton-chain σ has all its terminal and source states in $\Omega_{\text{final}}(\varphi)$ we shall call it a *skeleton-chain over φ* . We say that a finite sequence $\{x_0, x_1, \dots, x_m\}$ of points in Z is a σ -*chain over φ* iff it starts and finishes at points $x_0, x_m \in \Omega_{\text{final}}(\varphi)$ and for every $i = 1, \dots, m$, $x_i \in \sigma(x_{i-1})$. A skeleton-chain over φ can be seen as a finite set of σ -chains over φ which may cross each other, but without forming cycles. We will say that a skeleton-chain σ is subordinated to ψ if $\sigma(x) \subseteq \psi(x)$ for all $x \in Z$. We write then $\sigma \prec \psi$ to express the fact that

σ is subordinated to ψ . Given two skeleton-chains over φ , $\sigma_1 : Z_1 \rightarrow \mathcal{P}(Z_1)$ and $\sigma_2 : Z_2 \rightarrow \mathcal{P}(Z_2)$, we say that σ_1 is *epimorphic (over φ) to σ_2* iff there is a surjective map $h : Z_1 \rightarrow Z_2$ such that $h^{-1}(\sigma_2(x)) \subseteq \sigma_1(h^{-1}(x))$ for every $x \in Z_2$, and all states in $h^{-1}(x) \cup \{x\}$ belong to the same connected component of $\Omega_{\text{final}}(\varphi)$, for every terminal and source states $x \in Z_2$. This easily implies that every σ_2 -chain over φ can be lifted into a σ_1 -chain over φ with the same length, which starts and finishes exactly at the same components of $\Omega_{\text{final}}(\varphi)$.

Definition 6.1. Let $\mathcal{S} \subseteq \mathcal{C}^0(X)$ be a class of continuous mappings. We say that \mathcal{S} has the skeleton-chain perturbation property iff

1. \mathcal{S} is closed w.r.t. the metric d_H ,
2. Given $0 < r_0 < r$ there is $\epsilon > 0$ such that for every $f \in \mathcal{S}$ and any skeleton-chain σ over $f_{r_0}^\bullet$ subordinated to $f_{r+\epsilon}^\bullet$, there is some $g \in \mathcal{S}$ such that $f_{r_0}^\bullet \prec g_r^\bullet \prec f_{r+\epsilon}^\bullet$, and a skeleton-chain σ' over $f_{r_0}^\bullet$ epimorphic to σ which is subordinated to g_r^\bullet .

Theorem 6.2 is a corollary of the theorem and proposition bellow.

Theorem 6.3. Let $\mathcal{S} \subseteq \mathcal{C}^0(X)$ be a class of continuous mappings with the skeleton-chain perturbation property. Given $f \in \mathcal{S}$ and $\epsilon > 0$ there is some $g \in \mathcal{S}$ such that $g \prec f_\epsilon^\bullet$ and g is combinatorially equivalent to f_ϵ^\bullet . In particular, g has finitely many attractors.

Proposition 6.4. The class $\mathcal{S} = \mathcal{C}^0(X)$ has the skeleton-chain perturbation property.

Consider now any class $\mathcal{S} \subseteq \mathcal{C}^0(X)$ of continuous mappings with the skeleton-chain perturbation property and define

$$\mathcal{X} = \{ f_\epsilon^\bullet : f \in \mathcal{S}, \epsilon \geq 0 \}.$$

This space can be naturally identified with the product $\mathcal{S} \times [0, +\infty[$ through the correspondence $f_\epsilon^\bullet \equiv (f, \epsilon)$. With this identification in mind consider the projection mapping $\delta : \mathcal{X} \rightarrow [0, +\infty[$, defined by $\delta(f_\epsilon^\bullet) = \epsilon$. For each $\varphi \in \mathcal{X}$, $\delta(\varphi)$ will be called the *radius* of φ . Finally we define $\mathcal{X}^{\text{stable}}$ to be the set of all open maps $\varphi \in \mathcal{X}$ such that either $\delta(\varphi) = 0$ or else φ satisfies the combinatorial stability condition. Clearly, given any $f \in \mathcal{S}$, $f_\epsilon^\bullet \in \mathcal{X}^{\text{stable}}$ for all but a countable set of parameters $\epsilon > 0$.

Definition 6.2. A mapping $F : \mathcal{X}^{\text{stable}} \rightarrow \mathcal{X}^{\text{stable}}$ is called a compression operator iff $F(\varphi) = \varphi \Leftrightarrow \delta(\varphi) = 0$, and whenever $\delta(\varphi) > 0$

1. $F(\varphi) \prec \varphi$,
2. $0 < \delta(F(\varphi)) < \delta(\varphi)$, and
3. $F(\varphi) \bowtie \varphi$, i.e. $F(\varphi)$ and φ are combinatorially equivalent.

We shall denote by \mathcal{M} the set of all compression operators on $\mathcal{X}^{\text{stable}}$.

Proposition 6.5. $\mathcal{M} \neq \emptyset$, i.e. there are compression operators on $\mathcal{X}^{\text{stable}}$.

Proof. Given $\varphi \in \mathcal{X}^{\text{stable}}$ let $\epsilon(\varphi)$ be the greatest lower bound of all $\epsilon > 0$ given by proposition 5.17. Let then $\mathcal{G}(\varphi) = \{\psi \in \mathcal{X} : \varphi_\epsilon^\circ \prec \psi \prec \varphi \text{ with } \epsilon = \epsilon(\varphi)\}$. Clearly, any selection function $F : \mathcal{X}^{\text{stable}} \rightarrow \mathcal{X}$ with $F(\varphi) \in \mathcal{G}(\varphi)$ for all $\varphi \in \mathcal{X}^{\text{stable}}$, is a compression operator. Therefore, by the Axiom of Choice, there are compression operators. \square

Each compression operator $F \in \mathcal{M}$ will define a kind of dynamical system with discrete time modeled by the first uncountable ordinal Ω . For each ordinal $\alpha < \Omega$, the power F^α will be defined as a compression operator in \mathcal{M} . We shall see that radius function $\delta : \mathcal{X}^{\text{stable}} \rightarrow [0, +\infty[$ is some kind of Lyapounov function decreasing along orbits of F . More precisely, we shall see that, given any $\varphi \in \mathcal{X}^{\text{stable}}$, the sequence $\{\delta(F^\alpha(\varphi))\}_{\alpha < \Omega}$ decreases and is eventually constant equal to zero, which proves that $\{F^\alpha(\varphi)\}_{\alpha < \Omega}$ is eventually constant equal to some function $f \in \mathcal{S}$. It follows easily that $f \prec \varphi$ is combinatorially equivalent to φ . In particular f will have finitely many attractors. Let ω denote the first infinite ordinal, $\omega = \{0, 1, 2, \dots\} (\equiv \mathbb{N})$. Note that any ordinal $n < \omega$ is a natural number and so we can define F^n to be the usual n -fold composition $F^n = F \circ \dots \circ F$. The following proposition is the key to define these 'dynamical systems' and prove theorem 6.3. The assumption that \mathcal{S} has the skeleton-chain perturbation property is used in the proof of this proposition.

Proposition 6.6. Given $F \in \mathcal{M}$ there is another compression operator $F^\omega \in \mathcal{M}$ such that

$$\delta(F^\omega(\varphi)) \leq \inf_{n < \omega} \delta(F^n(\varphi)) .$$

This crucial fact will be proved in the end of this section. Referring to the power F^ω of an operator $F \in \mathcal{M}$, we now define recursively all powers F^α , where $\alpha < \Omega$ is any countable ordinal. More precisely we shall define operators $\text{Pow}_\alpha : \mathcal{M} \rightarrow \mathcal{M}$, one for each ordinal $\alpha < \Omega$.

Definition 6.3. For $n < \omega$, $\text{Pow}_n(F) = F^n$ is the usual n -fold composition. Pow_ω is an operator, that can be defined according to proposition 6.6. Given any $\omega < \alpha < \Omega$, if α is not a limit ordinal then $\alpha = \beta + 1$, and we define

$$\text{Pow}_\alpha(F) = F \circ \text{Pow}_\beta(F) = F \circ F^\beta .$$

Otherwise if α is a limit ordinal then $\alpha = \omega \cdot \beta = \overbrace{\omega + \omega + \dots}^{\beta \text{ times}}$ for some ordinal $\beta < \alpha$ and we define

$$\text{Pow}_\alpha(F) = \text{Pow}_\beta(\text{Pow}_\omega(F)) = (F^\omega)^\beta .$$

A sequence $\{r_\alpha\}_{\alpha < \Omega}$, of non-negative real numbers $r_\alpha \geq 0$, is said to be *strictly decreasing* iff $r_\alpha \leq \inf_{\beta < \alpha} r_\beta$ whenever α is a limit ordinal, and $r_{\alpha+1} < r_\alpha$ whenever $r_\alpha > 0$. The sequence $\{r_\alpha\}_{\alpha < \Omega}$ is said to be *eventually zero* iff there is some ordinal $\gamma < \Omega$ such that $r_\alpha = 0$ for all $\alpha \geq \gamma$.

Lemma 6.7. Every strictly decreasing sequence $\{r_\alpha\}_{\alpha < \Omega}$, of non-negative real numbers $r_\alpha \geq 0$, is eventually zero.

Proof. For each integer $n \geq 1$, the set $A_n = \{\alpha < \Omega : r_\alpha - r_{\alpha+1} \geq 1/n\}$ is finite. Therefore $\alpha_n = \cup A_n < \Omega$ is a countable ordinal. Since $\alpha_1 \leq \alpha_2 \leq \dots \alpha_n \leq \dots$ is a countable sequence of countable ordinals, $\gamma = \cup_{n=1}^\infty \alpha_n$ is again a countable ordinal. Clearly $r_\alpha = 0$ for every $\alpha \geq \gamma$. \square

Proposition 6.8. For every $\varphi \in \mathcal{X}^{\text{stable}}$ and $F \in \mathcal{M}$, the sequence $\{\delta(F^\alpha(\varphi))\}_{\alpha < \Omega}$ is strictly decreasing and, therefore, it is eventually zero.

Proof. Given $\alpha < \Omega$ such that $\delta(F^\alpha(\varphi)) > 0$, since $F^{\alpha+1}(\varphi) = F(F^\alpha(\varphi))$, by item 2. in the definition of a compression operator, it follows that $\delta(F^{\alpha+1}(\varphi)) > 0$. Let us prove, by transfinite induction in $\alpha < \Omega$, that

$$(P_\alpha) \text{ for all } \beta < \alpha, \varphi \in \mathcal{X}^{\text{stable}} \text{ and } F \in \mathcal{M}, \delta(F^\alpha(\varphi)) \leq \delta(F^\beta(\varphi)) .$$

If $\alpha = n$ is a finite ordinal it follows by definition 6.2 that a strict inequality holds. Proposition 6.6 shows that (P_ω) holds.

Assume now that (P_β) holds for all $\beta < \alpha$ and let us prove that (P_α) is also true. Suppose α is a limit ordinal. Otherwise (P_α) is trivial. Then there is some $\alpha_1 < \alpha$ such that $\alpha = \omega \cdot \alpha_1$. Now, given $\beta < \alpha$ there are $\beta_1 < \beta$ and $n < \omega$ such that $\beta = \omega \cdot \beta_1 + n$. To prove this define $\beta_2 = \min\{\beta' \leq \beta : \exists n < \omega \beta = \beta' + n\}$. Then $\beta = \beta_2 + n$ and β_2 is a limit ordinal. Therefore $\beta_2 = \omega \cdot \beta_1$ for some $\beta_1 < \beta_2 \leq \beta$. Since α is a limit ordinal $\omega \cdot \beta_1 + n = \beta < \alpha$ implies that $\omega \cdot (\beta_1 + 1) = \omega \cdot \beta_1 + \omega \leq \alpha = \omega \cdot \alpha_1$, which in turn implies that $\beta_1 + 1 \leq \alpha_1$. Thus

$$\begin{aligned} \delta(F^\alpha(\varphi)) &= \delta((F^\omega)^{\alpha_1}(\varphi)) \leq \delta((F^\omega)^{\beta_1+1}(\varphi)) \\ &= \delta(F^\omega((F^\omega)^{\beta_1}(\varphi))) \leq \delta(F^n(F^{\omega \cdot \beta_1}(\varphi))) = \delta(F^\beta(\varphi)) \end{aligned}$$

proving (P_α) . □

Proof. (of Theorem 6.3) Given $f \in \mathcal{S}$ and $\epsilon > 0$, taking ϵ smaller if necessary, we may assume that $\varphi = f_\epsilon^\bullet \in \mathcal{X}^{\text{stable}}$. Consider the transfinite sequence $\{F^\alpha(\varphi)\}_{\alpha < \Omega}$. Let $\gamma < \Omega$ be the first ordinal such that $\delta(F^\gamma(\varphi)) = 0$. Then $h = F^\gamma(\varphi) \in \mathcal{S}$ is a continuous map such that $h \prec \varphi$ and h is combinatorially equivalent to φ , which proves the theorem. □

Two easy lemmas are needed to prove proposition 6.6.

Lemma 6.9. $f_r^\bullet \prec g_s^\bullet \Leftrightarrow d_H(f, g) < s - r$.

Next lemma states that in an open map extension $\varphi_0 \prec \varphi_1$, the combinatorics of φ_1 missing in φ_0 can be encapsulated in some skeleton-chain σ over φ_0 which is subordinated to φ_1 .

Lemma 6.10. *Given open maps $\varphi_0, \varphi_1 \in \mathcal{O}_{\text{Cont}}(X)$ with $\varphi_0 \prec \varphi_1$, there is a skeleton-chain σ over φ_0 , subordinated to φ_1 , such that all open maps $\varphi_0 \leq \psi \leq \varphi_1$, admitting a skeleton-chain σ' over φ_0 epimorphic to σ , are combinatorially equivalent to φ_1 .*

Proof. (of proposition 6.6) Take any operator $F \in \mathcal{M}$. Given $\varphi \in \mathcal{X}^{\text{stable}}$, let $\varphi_n = (f_n)_{\delta_n}^\bullet$ be the sequence $\varphi_n = F^n(\varphi)$. The sequence $\{\delta_n\}$ is strictly decreasing and we set $r = \lim_{n \rightarrow \infty} \delta_n$. The sequence of maps $\{f_n\}$ is Cauchy because, by lemma 6.9 $(f_{n+1})_{\delta_{n+1}}^\bullet \prec (f_n)_{\delta_n}^\bullet$ implies $d_H(f_n, f_{n+1}) < \delta_n - \delta_{n+1}$, and the series $\sum_{n \geq 1} \delta_n - \delta_{n+1}$ is

summable. Since \mathcal{S} is complete we can take then $f \in \mathcal{S}$ such that $d_H(f_n, f) \rightarrow 0$ as $n \rightarrow \infty$. In order to define $F^\omega(\varphi)$ we consider two cases:

First case: $r = 0$. In this case we define $F^\omega(\varphi) = f$. Clearly $f \prec \varphi$ and f is combinatorially equivalent to φ .

Second case: $r > 0$. In this case take $0 < r_0 < r$, as in proposition 5.17, such that $f_{r_0}^\bullet$ is combinatorially equivalent to f_r^\bullet . Note that it may happen that f_r^\bullet is not combinatorially stable, and so not combinatorially equivalent to any φ_n . For the pair of positive real numbers $0 < r_0 < r$ we can, according to definition 6.1, take $\epsilon > 0$ in order to fulfill the second requirement of this definition. Then we have $\varphi_n \prec f_{r+\epsilon}^\bullet$, for all large enough n . Taking, if necessary, $\epsilon > 0$ smaller we can assume that $\varphi_n \prec f_{r+\epsilon}^\bullet \prec \varphi$. In particular these three open maps are combinatorially equivalent. Choose now a skeleton-chain σ over $f_{r_0}^\bullet$ subordinated to $f_{r+\epsilon}^\bullet$, as given in lemma 6.10. Then by definition of the 'skeleton-chain perturbation property', see 6.1, there is some map $g \in \mathcal{S}$ such that $f_{r_0}^\bullet \prec g_r^\bullet \prec f_{r+\epsilon}^\bullet$. There is also some chain skeleton σ' over $f_{r_0}^\bullet$, epimorphic to σ over $f_{r_0}^\bullet$, and subordinated to g_r^\bullet . Lemma 6.10 above now implies that g_r^\bullet is combinatorially equivalent to $f_{r+\epsilon}^\bullet$ and, therefore, to φ . In this case we define $F^\omega(\varphi) = g_r^\bullet$. \square

Finally to prove proposition 6.4 we shall use that

Lemma 6.11. *Given $f \in \mathcal{C}^0(X)$, $Z \subseteq X$ finite, and $\epsilon > \delta > 0$, there is some $g \in \mathcal{C}^0(X)$ with $d_H(f, g) < \epsilon$ such that $g(B_\delta(x)) \supseteq \overline{B_{\epsilon-\delta}(f(x))}$, $\forall x \in Z$.*

Proof. Take $0 < \eta < \delta$ so that the oscillation of f in each ball $B_\eta(x)$, with $x \in Z$, is less than $\delta > 0$. Then any map $g \in \mathcal{C}^0(X)$ which coincides with f outside $B_\eta(Z)$, and takes values in $\overline{B_{\epsilon-\delta}(f(x))}$ for arguments in $B_\eta(x)$, must satisfy $d_H(f, g) < \epsilon$. Of course we can choose such a g in a way that $g(B_\eta(x))$ covers $\overline{B_{\epsilon-\delta}(f(x))}$, for each $x \in Z$. \square

Proof. (of proposition 6.4) We only have to prove that $\mathcal{S} = \mathcal{C}^0(X)$ satisfies condition 2. in definition 6.1. Given $0 < r_0 < r$ take any $0 < \epsilon < r - r_0$. Take $f \in \mathcal{C}^0(X)$, and let $\sigma : Z \rightarrow \mathcal{P}(Z)$ be a skeleton-chain over $f_{r_0}^\bullet$ subordinated to $f_{r+\epsilon}^\bullet$. Choose then $0 < \delta < \epsilon/2$ such that:

- (a) $B_\delta(Z \cap \Omega_{\text{final}}(f_{r_0}^\bullet)) \subseteq \Omega_{\text{final}}(f_{r_0}^\bullet)$,
- (b) $\text{graph}(\sigma) \subseteq \text{graph}(f_{r+\epsilon-2\delta}^\bullet)$,

and take $g \in \mathcal{C}^0(X)$, according to lemma 6.11, such that $d_H(f, g) < \epsilon$ and, for all $x \in Z$, $g(B_\delta(x)) \supseteq \overline{B_{\epsilon-\delta}(f(x))}$. Since $d_H(f, g) < \epsilon < r - r_0$, by lemma 6.9, this implies that $f_{r_0}^\bullet \prec g_r^\bullet \prec f_{r+\epsilon}^\bullet$. Let us prove now that there is a skeleton-chain σ' over $f_{r_0}^\bullet$ epimorphic to σ and subordinated to g_r^\bullet . This will be proved by induction. To set up the induction scheme define $Z_0 = \{x \in Z : \sigma(x) = \emptyset\}$ and, for each $n \geq 1$, $Z_n = \{x \in Z : \sigma(x) \subseteq Z_{n-1}\}$. Clearly, for some $N \in \mathbb{N}$, $Z_0 \subset Z_1 \subset \dots \subset Z_N = Z$ and for each $0 \leq n \leq N$, $\sigma_n = \sigma|_{Z_n}$ is a skeleton-chain on Z_n . Define $Z'_0 = Z_0$, $h_0 : Z'_0 \rightarrow Z_0$ to be the identity map, and $\sigma'_0 = \sigma_0$. Assume, as our induction hypothesis, that there is a surjective map $h_{n-1} : Z'_{n-1} \rightarrow Z_{n-1}$ such that for all $x \in Z_{n-1}$, (1) $d(h_{n-1}^{-1}(x), x) < \delta$, and (2) $h_{n-1}^{-1}(\sigma(x)) = \sigma'_{n-1}(h_{n-1}^{-1}(x))$. Let us construct now Z'_n , σ'_n and $h_n : Z'_n \rightarrow Z_n$, surjective, satisfying the correspondent conditions (1') $d(h_n^{-1}(x), x) < \delta$, and (2') $h_n^{-1}(\sigma(x)) = \sigma'_n(h_n^{-1}(x))$, for every $x \in Z_n$. Take $x \in Z_n$ and let $\sigma(x) = \{y_1, \dots, y_m\} \subseteq Z_{n-1}$. For each $1 \leq i \leq m$, consider the pre-images $(h_{n-1})^{-1}(y_i) = \{y'_{i,1}, y'_{i,2}, \dots, y'_{i,k_i}\}$, with $k_i \geq 1$. Thus

$$(h_{n-1})^{-1}(\sigma(x)) = \{y'_{1,1}, \dots, y'_{1,k_1}, y'_{2,1}, \dots, y'_{2,k_2}, \dots, y'_{m,1}, \dots, y'_{m,k_m}\},$$

and, by item (1) of the induction hypothesis, each $y'_{i,j}$ in this set satisfies $d(y'_{i,j}, y_i) < \delta$. Therefore, since by item (b) above $y_i \in \sigma(x) \subseteq f_{r+\epsilon-2\delta}^\bullet(x)$,

$$\begin{aligned} d(y'_{i,j}, f(x)) &\leq d(y'_{i,j}, y_i) + d(y_i, f(x)) \\ &\leq \delta + r + \epsilon - 2\delta = r + \epsilon - \delta, \end{aligned}$$

and so

$$y'_{i,j} \in B_{r+\epsilon-\delta}(f(x)) = B_r(\overline{B_{\epsilon-\delta}(f(x))}) \subseteq B_r(g(B_\delta(x))) = g_r^\bullet(B_\delta(x)).$$

This shows that there is some $x'_{i,j} \in B_\delta(x)$ such that $y'_{i,j} \in g_r^\bullet(x'_{i,j})$. Clearly, this choice is not unique. In fact, there is an open set of possible choices for each $x'_{i,j}$. Therefore, we may assume that the choice of each $x'_{i,j}$, for every $x \in Z_n$, is made in a way that they all become distinct. Now, let $Z'_n(x) = \{x'_{i,j} : 1 \leq i \leq m, 1 \leq j \leq k_i\}$, and define $Z'_n = Z'_{n-1} \cup \bigcup_{x \in Z_n} Z'_n(x)$. Let $\sigma'_n : Z'_n \rightarrow \mathcal{P}(Z'_n)$ be the extension of σ'_{n-1} defined by $\sigma'_n(x'_{i,j}) = y'_{i,j}$, for every $x'_{i,j} \in Z'_n(x)$. Finally, define $h_n : Z'_n \rightarrow Z_n$ extending h_{n-1} in such a way that $(h_n)^{-1}(x) = Z'_n(x)$ for every $x \in Z_n \setminus Z_{n-1}$. By the remark above the sets $Z'_n(x)$ are pairwise disjoint, which shows that h_n is a well defined single valued map. By construction h_n is clearly surjective. We have

$$\sigma'_n(h_n^{-1}(x)) = \sigma'_n(Z'_n(x)) = \{y'_{i,j} : 1 \leq i \leq m, 1 \leq j \leq k_i\} = (h_{n-1})^{-1}(\sigma(x)),$$

which establishes (2'). On the other hand, since $x'_{i,j} \in B_\delta(x)$, $d(h_n(x'_{i,j}), x'_{i,j}) = d(x, x'_{i,j}) < \delta$, for every $x'_{i,j} \in Z'_n(x)$, which proves (1'). Thus, by induction, (1') and (2') hold for all n up to N . Let then $Z' = Z'_N$, $\sigma' = \sigma'_N$ and $h = h_N$. By item (1') of the just proved statement we have, for all $x \in Z$, $d(h^{-1}(x), x) < \delta$. Let $x \in Z$ be a terminal or source state of σ . Then $x \in \Omega_{\text{final}}(f_{r_0}^\bullet)$, and, by (a) above, $h^{-1}(x) \cup \{x\}$ is contained in a single connected component of $\Omega_{\text{final}}(f_{r_0}^\bullet)$. We have constructed a skeleton-chain σ' over $f_{r_0}^\bullet$ which is epimorphic to σ over $f_{r_0}^\bullet$, and subordinated to g_r^\bullet , thus proving item 2. of definition 6.1. \square

References

- [A] V. Araújo, *Attractors and time averages for random maps*, Ann. Inst. H. Poincaré Analyse non lin. 17, pp 307-369, 2000
- [AF] J-P. Aubin, H. Frankowska, *Set-Valued Analysis*, Birkhäuser, Boston, 2000
- [B] P. Billingsley, *Probability and Measure*, Wiley-Interscience, New York, 1985
- [F] K. J. Falconer, *The geometry of fractal sets*, Cambridge Tracts in Mathematics; 85, Cambridge University Press, 1985
- [HPS] M.W. Hirsh, C.C. Pugh, and M. Shub *Invariant manifolds*, Springer Lect. Notes in Math. 583 Berlin-New York, 1977
- [N] S. Nadler Jr. *Hyperspaces of sets*, Pure and Applied Mathematics, A Series of Monographs and Textbooks, Marcel Dekker Inc., 1978
- [P] J. Palis, *A global view of dynamics and a conjecture on the denseness of finitude of attractors*, Astérisque, 1998
- [Pi] K. Sergei Yu. Pilyugin, *The space of dynamical systems with the C^0 - topology*, Springer Lect. Notes in Math. 1571 Berlin-New York, 1994
- [S] M. Shub, *Global stability of dynamical systems*, Springer-Verlag, New York, 1987
- [Sm] S. Smale, *On gradient dynamical systems*, Ann. of Math. (2) 74, pp 199-206, 1961
- [T] F. Takens, *On Zeeman's tolerance stability conjecture*, Manifolds - Amsterdam, Lect. Notes in Math. 197, Springer-Verlag, pp 209-219, 1971
- [V1] M. Viana, *Stochastic dynamics of deterministic systems*, Lect. Notes XXI Braz. Math Colloq., IMPA, 1997
- [V2] M. Viana, *Dynamics: a probabilistic and geometric perspective*, Proceedings ICM, 1998 - Documenta Mathematica